

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶: C07K 16/24, 16/46, 17/02, C07H 15/12, G01N 33/53, C12P 21/08, C12N 5/10, 5/20, 15/13, A61K 39/395	A1	(11) International Publication Number: WO 95/07301 (43) International Publication Date: 16 March 1995 (16.03.95)
(21) International Application Number: PCT/US94/10308 (22) International Filing Date: 7 September 1994 (07.09.94) (30) Priority Data: 08/117,366 7 September 1993 (07.09.93) US 08/136,783 14 October 1993 (14.10.93) US (60) Parent Application or Grant (63) Related by Continuation US 08/136,783 (CIP) Filed on 14 October 1993 (14.10.93) (71) Applicants (for all designated States except US): SMITHK- LINE BEECHAM CORPORATION [US/US]; Corporate In- tellectual Property, UW2220, 709 Swedeland Road, P.O. Box 1539, King of Prussia, PA 19406-0939 (US). SMITHK- LINE BEECHAM PLC [GB/GB]; New Horizons Court, Brentford TW8 9BD (GB). (72) Inventors; and (75) Inventors/Applicants (for US only): HOLMES, Stephen, Dud- ley [GB/GB]; Yew Tree Bottom Road, Epsom, Surrey KT15 5XQ (GB). GROSS, Mitchell, Stuart [US/US]; 667 Pugh		<p>Road, Wayne, PA 19087 (US). SYLVESTER, Daniel, R. [US/US]; 42 Rossiter Avenue, Phoenixville, PA 19460 (US).</p> <p>(74) Agents: SUTTON, Jeffrey, A. et al.; SmithKline Beecham Corporation, Corporate Intellectual Property, UW2220, 709 Swedeland Road, P.O. Box 1539, King of Prussia, PA 19406-0939 (US).</p> <p>(81) Designated States: AM, AU, BB, BG, BR, BY, CA, CN, CZ, EE, FI, GE, HU, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LV, MD, MG, MN, MW, NO, NZ, PL, RO, RU, SD, SI, SK, TJ, TT, UA, US, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD).</p> <p>Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the</i> <i>claims and to be republished in the event of the receipt of</i> <i>amendments.</i></p>
(54) Title: RECOMBINANT IL4 ANTIBODIES USEFUL IN TREATMENT OF IL4 MEDIATED DISORDERS (57) Abstract Chimeric and humanized IL4 MAbs derived from high affinity MAbs, pharmaceutical compositions containing same, and methods of treatment are provided.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

RECOMBINANT IL4 ANTIBODIES USEFUL IN TREATMENT OF IL4
MEDIATED DISORDERS

Cross Reference to Related Applications

- 5 This application is a continuation-in-part of U.S. Serial No. 08/136,783 filed October 14, 1993 which is a continuation of U.S. Serial No. 08/117,366 filed September 7, 1993 both of which are hereby incorporated by reference in their entirety.

10 Field of the Invention

 The present invention relates generally to the field of fusion proteins, and to proteins useful in treatment and diagnosis of conditions mediated by IL4 and excess IgE production, and more specifically to chimeric and humanized IL4 antibodies.

15 Background of the Invention

- Atopic allergic diseases range from the relatively minor, such as seasonal rhinitis and conjunctivitis, to the more serious, such as atopic dermatitis and atopic asthma, and life threatening, such as anaphylactic shock. Linking these conditions is the immune response of the body to allergens, which response involves the
20 production of immunoglobulin E (IgE) antibodies in genetically predisposed individuals (atopy). Inhibition of IgE production has long been a goal in specific immunotherapy of allergic disease using desensitization vaccines. However, in recent years the safety and efficacy of vaccine therapy have been questioned, but the desire to reduce IgE levels has not waned.

- 25 Interleukin 4 (IL4) is a protein mediator in the lymphoid system. Studies of lymphocytes from atopic individuals have revealed the presence of higher than normal numbers of T lymphocytes with the ability to secrete IL4 in response to stimulation, and larger quantities of IL4 secreted following stimulation.

- Anti-IL4 antibody has been found to inhibit IgE, but not IgG₁ or IgG₂,
30 [Finkelman *et al*, Ann. Rev. Immunol., 8:303 (1990)], and the production of IL5 secreting T cells [Maggi *et al*, J. Immunol., 148:2142 (1992)]. Further, recent data suggests that IL4 may affect eosinophil accumulation in tissues. See, e.g. Tepper *et al*, Cell, 62:457 (1990); Tepper *et al*, Cell, 57:503 (1989).

- There remains a need in the art for a high affinity IL4 antagonist, which
35 would reduce eosinophil inflammation both by reducing the proliferation of IL5 secreting cells, and by inhibiting an adherence mechanism whereby eosinophils may be accumulating in tissues, and can be used to treat, prevent or diagnose allergic reactions.

Summary of the Invention

In a first aspect, the present invention provides a fusion protein having a binding affinity for human interleukin-4 which comprises complementarity determining regions (CDRs) derived from a non-human neutralizing monoclonal antibody (MAb) characterized by a dissociation constant equal to or less than 2×10^{-10} M for human IL4, and a first fusion partner in which at least one, and preferably all complementarity determining regions (CDRs) of the first fusion partner are replaced by CDRs from the non-human monoclonal antibody (MAb). The non-human neutralizing monoclonal antibody may be selected from the group consisting of 3B9 and 6A1 as described more fully in the Detailed Description. Preferably, the fusion protein is operatively linked to a second fusion protein as well, which comprises all or a part of an immunoglobulin constant chain.

In a related aspect, the present invention provides CDRs derived from non-human neutralizing monoclonal antibodies (MAb) characterized by a dissociation constant equal to or less than 2×10^{-10} M for human IL4, and nucleic acid molecules encoding such CDRs.

In another aspect, the invention provides humanized antibodies having at least one, and preferably six, complementarity determining regions (CDRs) derived from non-human neutralizing monoclonal antibodies (MAb) characterized by a dissociation constant equal to or less than 2×10^{-10} M for human IL4.

In still another aspect, there is provided a chimeric antibody containing human heavy and light chain constant regions and heavy and light chain variable regions derived from non-human neutralizing monoclonal antibodies (MAb) characterized by a dissociation constant equal to or less than 2×10^{-10} M for human IL4.

In still another aspect, the present invention provides a pharmaceutical composition which contains one (or more) of the above-described fusion proteins or MAbs (e.g., humanized, chimeric, etc.) and a pharmaceutically acceptable carrier.

In a further aspect, the present invention provides a method for treating and/or preventing allergic conditions in humans by administering to said human an effective amount of pharmaceutical composition of the invention.

In yet another aspect, the present invention provides methods for, and components useful in, the recombinant production of the fusion proteins, MAbs (e.g., humanized, chimeric, etc.), CDRs thereof, a Fab, or $F(ab)_2$, or analog thereof which is derived from non-human neutralizing monoclonal antibodies (MAb) characterized by a dissociation constant equal to or less than 2×10^{-10} M for human

IL4. These components include isolated nucleic acid sequences encoding same, recombinant plasmids containing the nucleic acid sequences under the control of selected regulatory sequences capable of directing the expression thereof in host cells, and host cells (preferably mammalian) transfected with the recombinant
5 plasmids. The production method involves culturing a transfected host cell line of the present invention under conditions such that an antibody, preferably a humanized antibody, is expressed in said cells and isolating the expressed product therefrom.

10 In yet another aspect of the invention is a method to diagnose allergies and other conditions associated with excess immunoglobulin E production in a human which comprises contacting a sample of biological fluid with the fusion proteins, MABs (e.g., humanized, chimeric, etc.) and Fabs of the instant invention and assaying for the occurrence of binding between said fusion protein, MAb or Fab and human interleukin 4.

15 In another related aspect is provided a method for screening monoclonal antibodies which have a high titer for human interleukin 4 which comprises: (a) preparing a hybridoma cell line characterized by secretion of a monoclonal antibody to human interleukin 4; and (b) screening said hybridoma cell line with aldehyde-coupled human interleukin-4 or biotinylated human interleukin-4. Preferably, the
20 hybridoma cell line is screened with biotinylated human interleukin-4.

Also provided is a neutralizing MAB having high affinity for IL4, a Fab fragment or a F(ab')₂ fragment thereof, produced by screening a library of hybridoma products with aldehyde-coupled human interleukin-4 or biotinylated human IL4.

25 In another aspect, the present invention provides rodent neutralizing monoclonal antibodies specific for human interleukin-4 and having a binding affinity characterized by a dissociation constant equal to or less than about 2×10^{-10} M. Exemplary of such monoclonal antibodies is the murine MAB, 3B9, and the rat MAB, 6A1 and other MABs have the same identifying characteristics (i.e., binds to
30 the same epitope(s) as 3B9 or 6A1 with a specificity for human IL4 and a dissociation constant equal to or less than about 2×10^{-10} M). Another aspect of the invention is hybridoma 3426A11C1B9.

Other aspects and advantages of the present invention are described further in the following detailed description of the preferred embodiments thereof.

Brief Description of the Drawings

Fig. 1 [SEQ ID NOS: 1 and 2] illustrates the light chain variable region (amino acids 21-132) for the murine IL4 antibody 3B9, and the human/murine 3B9 chimeric antibody as well as the native signal sequence (amino acids 1-20). The underlined portions indicate the CDRs [SEQ ID NOS: 15 and 16; SEQ ID NOS: 17 and 18; and SEQ ID NOS: 19 and 20].

Fig. 2 [SEQ ID NOS: 3 and 4] illustrates the heavy chain variable region (amino acids 20-140) of the murine 3B9, and the native signal sequence (amino acids 1-19). The underlined portions indicate the CDRs [SEQ ID NOS: 21 and 22; SEQ ID NOS: 23 and 24; and SEQ ID NOS: 25 and 26].

Fig. 3 [SEQ ID NOS: 9 and 10] illustrates the heavy chain variable region (amino acids 21-141) of the human/murine 3B9 chimeric antibody and its signal sequence (amino acids 1-19; SEQ ID NOS: 5 and 6). The underlined portions indicate the CDRs derived from 3B9 [SEQ ID NOS: 21 and 22; SEQ ID NOS: 23 and 24; and SEQ ID NOS: 25 and 26].

Fig. 4 [SEQ ID NOS: 11 and 12] illustrates the heavy chain variable region (amino acids 20-141) of the humanized 3B9 antibody and a signal sequence (amino acids 1-19; SEQ ID NOS: 5 and 6). The underlined portions indicate the CDRs derived from 3B9 [SEQ ID NOS: 54 and 22; SEQ ID NOS: 55 and 24; and SEQ ID NOS: 56 and 26].

Fig. 5 [SEQ ID NOS: 13 and 14] illustrates the light chain variable region (amino acids 21-131) of the humanized 3B9 antibody and a signal sequence (amino acids 1-20; SEQ ID NOS: 7 and 8). The underlined portions indicate the CDRs derived from 3B9 [SEQ ID NOS: 53 and 16; SEQ ID NOS: 17 and 18; and SEQ ID NOS: 27 and 28].

Fig. 6A [SEQ ID NOS: 5 and 6] is a heavy chain signal sequence used in Example 4 below.

Fig. 6B [SEQ ID NOS: 7 and 8] is a light chain signal sequence used in Example 4 below.

Fig. 7 is a schematic drawing of plasmid pIL4chhc3-pcd employed to express a chimeric IL4 heavy chain in mammalian cells. The plasmid contains a beta lactamase gene (BETA LAC), an SV-40 origin of replication (SV40), a cytomegalovirus promoter sequence (CMV), a signal sequence, the chimeric variable heavy chain of SEQ ID NOS: 9 and 10, a human heavy chain constant region, a poly A signal from bovine growth hormone (BGH), a betaglobin promoter (beta glopro), a dihydrofolate reductase gene (DHFR), and another BGH sequence poly A signal in a pUC19 background.

Fig. 8 is a schematic drawing of plasmid pIL4chlc-pcdn employed to express the chimeric IL4 light chain variable region of SEQ ID NOS: 1 and 2 in mammalian cells. The plasmid differs from that of Fig. 7 by containing a chimeric light chain variable region rather than that of the chimeric heavy chain, a human light chain constant region and a neomycin gene (Neo) in addition to DHFR.

Fig. 9 is a schematic drawing of plasmid pIL4hzhc-1-pcd employed to express the synthetic IL4 heavy chain variable region of SEQ ID NOS: 11 and 12 in mammalian cells. The plasmid differs from that of Fig. 7 by containing a humanized heavy chain variable region rather than that of the chimeric heavy chain.

Fig. 10 is a schematic drawing of plasmid pIL4hzlc1-0-Pcn employed to express the humanized IL4 light chain variable region of SEQ ID NOS: 13 and 14 in mammalian cells. The plasmid differs from that of Fig. 8 by containing a humanized light chain variable region rather than that of the chimeric light chain and does not encode the DHFR gene.

Detailed Description of the Invention

The present invention provides a variety of antibodies, fragments thereof, and fusion proteins particularly humanized antibodies, which are characterized by human IL4 binding specificity, neutralizing activity, and high affinity for human IL4 as exemplified in murine MA b 3B9 or the rat MA b 6A1. These products are useful in therapeutic and pharmaceutical compositions for treating IL4-mediated and IgE-mediated allergic reactions. These products are also useful in the diagnosis of an IL4 mediated condition by measurement (e.g., by enzyme linked immunosorbent assay (ELISA)) of circulating, endogenous IL4 levels in humans.

1. Definitions.

"Fusion protein" refers to a protein encoded by a fusion molecule, which may be obtained by expression in a selected host cell. Such fusion proteins are engineered antibodies, e.g., chimeric or humanized antibodies, or antibody fragments lacking all or part of an immunoglobulin constant region, e.g., Fv, Fab, or F(ab)₂ and the like.

"Fusion molecule" refers to a nucleic acid sequence encoding the complementarity determining regions (CDRs) from a non-human immunoglobulin that are inserted into a first fusion partner comprising human variable framework sequences. Optionally, the first fusion partner is operatively linked to a second fusion partner.

"First fusion partner" refers to a nucleic acid sequence encoding a human framework or human immunoglobulin variable region in which the native (or

naturally-occurring) CDRs are replaced by the CDRs of a donor antibody. The human variable region can be an immunoglobulin heavy chain, a light chain (or both chains), an analog or functional fragments thereof. Such CDRs or CDR regions, located within the variable region of antibodies (immunoglobulins) can be
5 determined by known methods in the art. For example Kabat et al., [Sequences of Proteins of Immunological Interest, 4th Ed., U.S. Department of Health and Human Services, National Institutes of Health (1987)], disclose rules for locating CDRs. In addition, computer programs are known which are useful for identifying CDR regions/structures.

10 The term "high titer" refers to an antibody having a binding affinity characterized by a K_d equal to or less than 2×10^{-10} M for human IL4.

By "binding specificity for human IL4" is meant a high titer (or affinity) for human, not bovine or murine, IL4.

"Second fusion partner" refers to another nucleotide sequence encoding a
15 protein or peptide to which the first fusion partner is fused in frame or by means of an optional conventional linker sequence (i.e., operatively linked). Preferably it is an immunoglobulin. The second fusion partner may include a nucleic acid sequence encoding the entire constant region for the same (i.e., homologous - the first and second fusion proteins are derived from the same source) or an additional (i.e.,
20 heterologous) antibody of interest. It may be an immunoglobulin heavy chain or light chain (or both chains as part of a single polypeptide). The second fusion partner is not limited to a particular immunoglobulin class or isotype. In addition, the second fusion partner may comprise part of an immunoglobulin constant region, such as found in a Fab, or $F(ab)_2$ (i.e., a discrete part of an appropriate human
25 constant region or framework region). Such second fusion partner may also comprise a sequence encoding an integral membrane protein exposed on the outer surface of a host cell, e.g., as part of a phage display library, or a sequence encoding a protein for analytical or diagnostic detection, e.g., horseradish peroxidase, β -galactosidase, etc.

30 The terms Fv, Fc, Fab, or $F(ab)_2$ are used with their standard meanings (see, e.g., Harlow et al., Antibodies A Laboratory Manual, Cold Spring Harbor Laboratory, (1988)).

As used herein, an "engineered antibody" describes a type of fusion protein, i.e., a synthetic antibody (e.g., a chimeric or humanized antibody) in which
35 portion of the light and/or heavy chain variable domains of a selected acceptor antibody are replaced by analogous parts from one or more donor antibodies which have specificity for the selected epitope. For example, such molecules may include

antibodies characterized by a humanized heavy chain associated with an unmodified light chain (or chimeric light chain), or vice versa. Engineered antibodies may also be characterized by alteration of the nucleic acid sequences encoding the acceptor antibody light and/or heavy variable domain framework regions in order to retain donor antibody binding specificity. These antibodies can comprise replacement of one or more CDRs (preferably all) from the acceptor antibody with CDRs from a donor antibody described herein.

A "chimeric antibody" refers to a type of engineered antibody which contains naturally-occurring variable region (light chain and heavy chains) derived from a donor antibody in association with light and heavy chain constant regions derived from an acceptor antibody.

A "humanized antibody" refers to a type of engineered antibody having its CDRs derived from a non-human donor immunoglobulin, the remaining immunoglobulin-derived parts of the molecule being derived from one (or more) human immunoglobulin. In addition, framework support residues may be altered to preserve binding affinity (see, e.g., Queen et al., Proc. Natl Acad Sci USA, 86:10029-10032 (1989), Hodgson et al., Bio/Technology, 9:421 (1991)).

The term "donor antibody" refers to an antibody (polyclonal, monoclonal, or recombinant) which contributes the nucleic acid sequences of its variable regions, CDRs, or other functional fragments or analogs thereof to a first fusion partner, so as to provide the fusion molecule and resulting expressed fusion protein with the antigenic specificity and neutralizing activity characteristic of the donor antibody. One donor antibody suitable for use in this invention is a non-human neutralizing monoclonal antibody (i.e., murine) designated as 3B9. The antibody 3B9 is defined as a high titer, human-IL4 specific (i.e., does not recognize bovine or murine IL4), neutralizing antibody of isotype IgG₁ having the variable light chain DNA and amino acid sequences of SEQ ID NOS: 1 and 2, and the variable heavy chain DNA and amino acid sequences of SEQ ID NOS: 3 and 4 on a suitable murine IgG constant region.

The term "acceptor antibody" refers to an antibody (polyclonal, monoclonal, or recombinant) heterologous to the donor antibody, which contributes all (or any portion, but preferably all) of the nucleic acid sequences encoding its heavy and/or light chain framework regions and/or its heavy and/or light chain constant regions to the second fusion partner. Preferably a human antibody is the acceptor antibody.

"CDRs" are defined as the complementarity determining region amino acid sequences of an antibody which are the hypervariable regions of immunoglobulin heavy and light chains. See, e.g., Kabat et al., Sequences of Proteins of

Immunological Interest, 4th Ed., U.S. Department of Health and Human Services, National Institutes of Health (1987). There are three heavy chain and three light chain CDRs (or CDR regions) in the variable portion of an immunoglobulin. Thus, "CDRs" as used herein refers to all three heavy chain CDRs, or all three light chain CDRs (or both all heavy and all light chain CDRs, if appropriate).

CDRs provide the majority of contact residues for the binding of the antibody to the antigen or epitope. CDRs of interest in this invention are derived from donor antibody variable heavy and light chain sequences, and include analogs of the naturally occurring CDRs, which analogs also share or retain the same antigen binding specificity and/or neutralizing ability as the donor antibody from which they were derived.

By 'sharing the antigen binding specificity or neutralizing ability' is meant, for example, that although MAb 3B9 may be characterized by a certain level of antigen affinity, and a CDR encoded by a nucleic acid sequence of 3B9 in an appropriate structural environment may have a lower or higher affinity, it is expected that CDRs of 3B9 in such environments will nevertheless recognize the same epitope(s) as 3B9. Exemplary heavy chain CDRs of 3B9 include SEQ ID NO: 22; SEQ ID NO: 24; SEQ ID NO: 26; and exemplary light chain CDRs of 3B9 include SEQ ID NO: 16; SEQ ID NO: 18; and SEQ ID NO: 20.

A "functional fragment" is a partial heavy or light chain variable sequence (e.g., minor deletions at the amino or carboxy terminus of the immunoglobulin variable region) which retains the same antigen binding specificity and/or neutralizing ability as the antibody from which the fragment was derived.

An "analog" is an amino acid sequence modified by at least one amino acid, wherein said modification can be chemical or a substitution or a rearrangement of a few amino acids (i.e., no more than 10), which modification permits the amino acid sequence to retain the biological characteristics, e.g., antigen specificity and high titer or affinity, of the unmodified sequence. For example, silent mutations can be constructed, via substitutions, to create endonuclease restriction sites within or surrounding CDR regions.

Analogous may also arise as allelic variations. An "allelic variation or modification" is an alteration in the nucleic acid sequence encoding the amino acid or peptide sequences of the invention. Such variations or modifications may be due to degeneracies in the genetic code or may be deliberately engineered to provide desired characteristics. These variations or modifications may or may not result in alterations in any encoded amino acid sequence. For example, the amino acid sequences of the light chain CDR SEQ ID NO: 16 are identical for the native

murine and humanized 3B9 antibody. However, this CDR sequence is encoded by both SEQ ID NO: 15 and SEQ ID NO: 53. Similarly, CDR SEQ ID NO: 22 is encoded both by SEQ ID NO: 21 and SEQ ID NO: 54; CDR SEQ ID NO: 24 is encoded both by SEQ ID NO: 23 and SEQ ID NO: 55; and CDR SEQ ID NO: 26 is encoded both by SEQ ID NO: 25 and SEQ ID NO: 56.

The term "effector agents" refers to non-protein carrier molecules to which the fusion proteins, and/or natural or synthetic light or heavy chain of the donor antibody or other fragments of the donor antibody may be associated by conventional means. Such non-protein carriers can include conventional carriers used in the diagnostic field, e.g., polystyrene or other plastic beads, polysaccharides, e.g., as used in the BIAcore [Pharmacia] system, or other non-protein substances useful in the medical field and safe for administration to humans and animals. Other effector agents may include a macrocycle, for chelating a heavy metal atom, or radioisotopes. Such effector agents may also be useful to increase the half-life of the fusion proteins, e.g., polyethylene glycol.

II. High Affinity IL4 Monoclonal Antibodies

For use in constructing the antibodies, fragments and fusion proteins of this invention, a non-human species (for example, bovine, ovine, primate, rodent (e.g., murine and rat), etc.) may be employed to generate a desirable immunoglobulin upon presentment with native human IL4 or a peptide epitope therefrom. Conventional hybridoma techniques are employed to provide a hybridoma cell line secreting a non-human MAb to IL4. Such hybridomas are then screened using IL4 covalently attached to 96-well plates or alternatively with biotinylated IL4 for use in a screening assay, as described in detail in Example 2 below. Thus one feature of the instant invention is a method to detect MAbs for human IL4 in which the assay systems avoid denaturing of IL4. In such a manner, it was discovered that high titer (or high affinity) MAbs to human IL4 can be detected.

As one example, the production of a high titer, neutralizing MAb from a murine donor is disclosed for the first time. MAb 3B9, which is a desirable murine (donor) antibody for use in developing a chimeric or humanized antibody, is described in detail in Example 1 below. The 3B9 MAb is characterized by an antigen binding specificity for human IL4, with a K_d of less than 2.0×10^{-10} M (about 1.8×10^{-10} M) for IL4. The K_d for IL4 of a Fab fragment of this 3B9 is less than about 3×10^{-10} M. The epitope of this antibody could not be mapped to IL4 with linear peptides, and hence the epitope is considered to bind to a non-contiguous epitope. The pattern of binding suggests a binding site at the B-C loop (residues 60-69) \rightarrow C helix (residues 70-93) region. These regions refer to the map designations

provided in Cook et al, J. Mol. Biol., 218:675-678 (1991), Walter et al, J. Biol. Chem., 267:20371-20376 (1992), Wlodaver et al, FEBS Lett., 309:59-64 (1992), Redfield et al, Biochem., 30:11029-11035 (1991), Smith et al, J. Mol. Biol., 224:899-904 (1992), Garrett et al, (1992), and Powers et al, Biochem., 31:4334-4346 (1992) and Science, 256:1673-1677 (1992), incorporated by reference herein.

Another desirable donor antibody is the rat MAb, 6A1. The production of this MAb is provided below in Example 7. This MAb is characterized by being isotype IgG_{2a}, and having a dissociation constant for hIL4 of less than 2.0×10^{-10} M (about 1.6×10^{-10} M). As with 3B9, the target epitope of this 6A1 does not map with IL4 linear peptides, and the epitope is therefore considered to be non-contiguous and three dimensional. The pattern of binding to IL4 muteins and its biological activity indicates binding in the D helix region of human IL4 (amino acid residues 109-127), most likely around the tyrosine at amino acid residue #124.

This invention is not limited to the use of the 3B9 MAb, the 6A1 MAb, or its hypervariable (i.e., CDR) sequences. Any other appropriate high titer IL4 antibodies characterized by a dissociation constant equal or less than 2.0×10^{-10} M for human IL4 and corresponding anti-IL4 CDRs may be substituted therefor. Wherever in the following description the donor antibody is identified as 3B9 or 6A1, this designation is made for illustration and simplicity of description only.

III. Antibody Fragments

The present invention also includes the use of Fab fragments or F(ab')₂ fragments derived from MAbs directed against human IL4. These fragments are useful as agents protective *in vivo* against IL4- and IgE- mediated conditions or *in vitro* as part of an IL4 diagnostic. A Fab fragment contains the entire light chain and amino terminal portion of the heavy chain; and an F(ab')₂ fragment is the fragment formed by two Fab fragments bound by disulfide bonds. MAbs 3B9, 6A1, and other similar high affinity, IL4 binding antibodies, provide sources of Fab fragments and F(ab')₂ fragments which can be obtained by conventional means, e.g., cleavage of the MAb with the appropriate proteolytic enzymes, papain and/or pepsin, or by recombinant methods. These Fab and F(ab')₂ fragments are useful themselves as therapeutic, prophylactic or diagnostic agents, and as donors of sequences including the variable regions and CDR sequences useful in the formation of recombinant or humanized antibodies as described herein.

IV. Anti-IL4 Amino Acid and Nucleotide Sequences of Interest

The MAb 3B9 or other antibodies described above may contribute sequences, such as variable heavy and/or light chain peptide sequences, framework sequences, CDR sequences, functional fragments, and analogs thereof, and the

nucleic acid sequences encoding them, useful in designing and obtaining various fusion proteins (including engineered antibodies) which are characterized by the antigen binding specificity of the donor antibody.

As one example, the present invention thus provides variable light chain and
5 variable heavy chain sequences from the IL4 murine antibody 3B9 and sequences derived therefrom. The heavy chain variable region of 3B9 is characterized by amino acid residues 20 to 140 of SEQ ID NO: 4. The CDR regions are indicated by underlining in Fig. 2 and are provided in SEQ ID NO: 22; SEQ ID NO: 24; and
10 SEQ ID NO: 26. The light chain clone variable region of 3B9 is characterized by amino acid residues 21 to 132 of Fig. 1 [SEQ ID NO: 2]. The CDR regions are from amino acid residues 44-58 [SEQ ID NO: 16]; 74-80 [SEQ ID NO: 18]; and 113-121 [SEQ ID NO: 20].

Chimeric heavy chain variable region and signal nucleotide and amino acid sequences are provided. These sequences are identical to the 3B9 heavy chain with
15 the exception of the signal sequence. The chimeric heavy chain signal sequence is provided in SEQ ID NOS: 5 and 6. The CDR regions are indicated by underlining in Fig. 3 and are identical in amino acid sequence to the native murine CDRs [SEQ ID NOS: 21-26]. The chimeric light chain variable region nucleotide and amino acid sequences are identical to the unmodified 3B9 sequences (amino acid residues
20 21-132 of SEQ ID NO: 2), making use of the natural mouse signal sequences (amino acid residues 1-20 of SEQ ID NO: 2).

A humanized heavy chain variable region and signal sequences are illustrated in Fig. 4 [SEQ ID NO: 11 and 12]. The signal sequence is also provided
25 in SEQ ID NO: 5 and 6. Other suitable signal sequences, known to those of skill in the art, may be substituted for the signal sequences exemplified herein. The CDR amino acid sequences of this construct are identical to the native murine and chimeric heavy chain CDRs and are provided by SEQ ID NO: 22 (encoded by SEQ ID NO: 54), SEQ ID NO: 24 (encoded by SEQ ID NO: 55), and SEQ ID NO: 56 (encodes SEQ ID NO: 26).

30 An exemplary (synthetic) humanized light chain variable sequence is illustrated in Fig. 5 [SEQ ID NOS: 13 and 14]. The signal sequence spans amino acid residues 1 to 19 of SEQ ID NO: 8. The CDR sequences of this figure are designated by underlining and differ from the CDR of the native murine CDR by a single amino acid of SEQ ID NO: 20. Thus, the CDRs of the humanized light chain
35 are provided by SEQ ID NO: 53 and 16, SEQ ID NO: 17 and 18, and SEQ ID NO: 27 and 28. This difference is described in detail in Example 3.

The nucleic acid sequences of this invention, or fragments thereof, encoding the variable light chain and heavy chain peptide sequences are used in unmodified form or are synthesized to introduce desirable modifications, e.g., restriction sites. The isolated naturally-occurring or alternatively synthetic nucleic acid sequences, which are derived from MAb 3B9 or from other desired high titer IL4 antibodies may optionally contain restriction sites to facilitate insertion or ligation into a suitable nucleic acid sequence such as encoding a desired antibody framework region, ligation with mutagenized CDRs or fusion with a nucleic acid sequence encoding a selected second fusion partner.

Taking into account the degeneracy of the genetic code, various coding sequences may be constructed which encode the variable heavy and light chain amino acid sequences, and CDR sequences of the invention as well as functional fragments and analogs thereof which share the antigen specificity of the donor antibody. The isolated nucleic acid sequences of this invention, or fragments thereof, encoding the variable chain peptide sequences or CDRs can be used to produce fusion proteins, chimeric or humanized antibodies, or other engineered antibodies of this invention when operatively combined with a second fusion partner.

These sequences are also useful for mutagenic introduction of specific changes within the nucleic acid sequences encoding the CDRs or framework regions, and for incorporation of the resulting modified or fusion nucleic acid sequence into a plasmid for expression. For example, silent substitutions in the nucleotide sequence of the framework and CDR-encoding regions were used to create restriction enzyme sites which facilitated insertion of mutagenized CDR (and/or framework) regions. These CDR regions were used in the construction of a humanized antibody of this invention.

It should be noted that in addition to isolated nucleic acid sequences encoding portions of the fusion protein and antibodies described herein, other such nucleic acid sequences may be employed, such as those complementary to the native sequences. Useful DNA sequences include those sequences which hybridize under stringent hybridization conditions [see, T. Maniatis et al, Molecular Cloning (A Laboratory Manual), Cold Spring Harbor Laboratory (1982), pages 387 to 389] to the DNA sequences. An example of one such stringent hybridization condition is hybridization at 4XSSC at 65°C, followed by a washing in 0.1XSSC at 65°C for an hour. Alternatively an exemplary stringent hybridization condition is in 50% formamide, 4XSSC at 42°C. Preferably, these hybridizing DNA sequences are at least about 18 nucleotides in length, i.e., about the size of a CDR.

V. *Fusion Molecules and Fusion Proteins*

Fusion molecules can encode fusion proteins which includes engineered antibodies such as, chimeric antibodies, and humanized antibodies. A desired fusion molecule contains CDR sequences encoding peptides having the antigen specificity of an IL4 antibody, preferably a high affinity antibody such as is provided by the present invention inserted into a first fusion partner (a human framework or human immunoglobulin variable region).

Preferably, the first fusion partner is operatively linked to a second fusion partner. The second fusion partner is defined above, and may include a sequence encoding a second antibody region of interest, for example an Fc region. Second fusion partners may also include sequences encoding another immunoglobulins to which the light or heavy chain constant region is fused in frame or by means of a linker sequence. Engineered antibodies directed against functional fragments or analogs of IL4 may be designed to elicit enhanced binding with the same antibody.

The second fusion partner may also be associated with effector agents as defined above, including non-protein carrier molecules, to which the second fusion partner may be operatively linked by conventional means.

Fusion or linkage between the second fusion partners, e.g., antibody sequences, and the effector agent may be by any suitable means, e.g., by conventional covalent or ionic bonds, protein fusions, or hetero-bifunctional cross-linkers, e.g., carbodiimide, glutaraldehyde, and the like. Such techniques are known in the art and readily described in conventional chemistry and biochemistry texts.

Additionally, conventional linker sequences which simply provide for a desired amount of space between the second fusion partner and the effector agent may also be constructed into the fusion molecule. The design of such linkers is well known to those of skill in the art.

In addition, signal sequences for the molecules of the invention may be modified to enhance expression. As one example a desired fusion protein having an amino acid sequence of the murine heavy chain sequence, which is identical to the chimeric variable heavy chain (V_H) of Fig. 2 [SEQ ID NO: 4], had the original signal peptide replaced with another signal sequence (amino acid residues 1-20) [SEQ ID NO: 6].

An exemplary fusion protein contains a variable heavy and/or light chain peptide or protein sequence having the antigen specificity of MAb 3B9, e.g., the V_H [amino acid residues 21-141 of SEQ ID NO: 9 and 10] and V_L chains [amino acid residues 21-132 of SEQ ID NOS: 1 and 2]. Still another desirable fusion protein of this invention is characterized by the amino acid sequence containing at least one,

and preferably all of the CDRs of the variable region of the heavy and/or light chains of the murine antibody molecule 3B9 with the remaining sequences being derived from a human source, or a functional fragment or analog thereof. See, e.g., the humanized V_H and V_L regions of SEQ ID NOS: 11 and 12 and SEQ ID NOS: 13 and 14 (Figs. 4 and 5).

In still a further embodiment, the engineered antibody of the invention may have attached to it an additional agent. For example, the procedure of recombinant DNA technology may be used to produce an engineered antibody of the invention in which the Fc fragment or CH3 domain of a complete antibody molecule has been replaced by an enzyme or other detectable molecule. (ie., a polypeptide effector or reporter molecule)

The second fusion partner may also be operatively linked to a non-immunoglobulin peptide, protein or fragment thereof heterologous to the CDR-containing sequence having the antigen specificity of murine 3B9. The resulting protein may exhibit both anti-IL4 antigen specificity and characteristics of the non-immunoglobulin upon expression. That fusion partner characteristic may be, e.g., a functional characteristic such as another binding or receptor domain, or a therapeutic characteristic if the fusion partner is itself a therapeutic protein, or additional antigenic characteristics.

Another desirable protein of this invention may comprise a complete antibody molecule, having full length heavy and light chains, or any discrete fragment thereof, such as the Fab or F(ab')₂ fragments, a heavy chain dimer, or any minimal recombinant fragments thereof such as an F_v or a single-chain antibody (SCA) or any other molecule with the same specificity as the selected donor MAb, e.g., MAb.3B9 or 6A1. Such protein may be used in the form of a fusion protein, or may be used in its unfused form.

Whenever the second fusion partner is derived from another antibody, e.g., any isotype or class of immunoglobulin framework or constant region, an engineered antibody results. Engineered antibodies can comprise immunoglobulin (Ig) constant regions and variable framework regions from one source, e.g., the acceptor antibody, and one or more (preferably all) CDRs from the donor antibody, e.g., the anti-IL4 antibody described herein. In addition, alterations, e.g., deletions, substitutions, or additions, of the acceptor MAb light and/or heavy variable domain framework region at the nucleic acid or amino acid levels, or the donor CDR regions may be made in order to retain donor antibody antigen binding specificity.

Such engineered antibodies are designed to employ one (or both) of the variable heavy and/or light chains of the IL4 MAb (optionally modified as

described) or one or more of the below-identified heavy or light chain CDRs (see Example 3). The engineered antibodies of the invention are neutralizing, i.e., they desirably block binding to the receptor of the IL4 protein. For example, the engineered antibody derived from MAb 3B9 is directed against a specific tertiary protein epitope of human IL4 believed to be at the B-C loop → C helix region, as described above.

Such engineered antibodies may include a humanized antibody containing the framework regions of a selected human immunoglobulin or subtype, or a chimeric antibody containing the human heavy and light chain constant regions fused to the IL4 antibody functional fragments. A suitable human (or other animal) acceptor antibody may be one selected from a conventional database, e.g., the KABAT® database, Los Alamos database, and Swiss Protein database, by homology to the nucleotide and amino acid sequences of the donor antibody. A human antibody characterized by a homology to the framework regions of the donor antibody (on an amino acid basis) may be suitable to provide a heavy chain constant region and/or a heavy chain variable framework region for insertion of the donor CDRs. A suitable acceptor antibody capable of donating light chain constant or variable framework regions may be selected in a similar manner. It should be noted that the acceptor antibody heavy and light chains are not required to originate from the same acceptor antibody.

Desirably the heterologous framework and constant regions are selected from human immunoglobulin classes and isotypes, such as IgG (subtypes 1 through 4), IgM, IgA, and IgE. However, the acceptor antibody need not comprise only human immunoglobulin protein sequences. For instance a gene may be constructed in which a DNA sequence encoding part of a human immunoglobulin chain is fused to a DNA sequence encoding a non-immunoglobulin amino acid sequence such as a polypeptide effector or reporter molecule.

One example of a particularly desirable humanized antibody contains CDRs of 3B9 inserted onto the framework regions of a selected human antibody sequence. For neutralizing humanized antibodies one, two or preferably three CDRs from the IL4 antibody heavy chain and/or light chain variable regions are inserted into the framework regions of the selected human antibody sequence, replacing the native CDRs of the latter antibody.

Preferably, in a humanized antibody, the variable domains in both human heavy and light chains have been engineered by one or more CDR replacements. It is possible to use all six CDRs, or various combinations of less than the six CDRs. Preferably all six CDRs are replaced. It is possible to replace the CDRs only in the

human heavy chain, using as light chain the unmodified light chain from the human acceptor antibody. Still alternatively, a compatible light chain may be selected from another human antibody by recourse to the conventional antibody databases. The remainder of the engineered antibody may be derived from any suitable acceptor human immunoglobulin.

The engineered humanized antibody thus preferably has the structure of a natural human antibody or a fragment thereof, and possesses the combination of properties required for effective therapeutic use, e.g., treatment of IL4 mediated inflammatory diseases in man, or for diagnostic uses.

As another example, an engineered antibody may contain three CDRs of the variable light chain region of 3B9 [SEQ ID NO: 16, 18, 20 and 28] and three CDRs of the variable heavy chain region of 3B9 [SEQ ID NO: 22, 24 and 26]. The resulting humanized antibody is characterized by the antigen binding specificity and high affinity of MAb 3B9.

It will be understood by those skilled in the art that an engineered antibody may be further modified by changes in variable domain amino acids without necessarily affecting the specificity and high affinity of the donor antibody (i.e., an analog). For example, humanized monoclonal antibodies have been constructed wherein the light chain amino acid residue at position 120 was an arginine [SEQ ID NO:13 and 14] or threonine [SEQ ID NOS:57 and 58]. It is anticipated that heavy and light chain amino acids may be substituted by other amino acids either in the variable domain frameworks or CDRs or both.

In addition, the constant region may be altered to enhance or decrease selective properties of the molecules of the instant invention. For example, dimerization, binding to Fc receptors, or the ability to bind and activate complement (see, e.g., Angal et al., Mol. Immunol., 30:105-108 (1993), Xu et al., J. Biol. Chem., 269:3469-3474 (1994), Winter et al., EP 307,434-B).

A fusion protein which is a chimeric antibody differs from the humanized antibodies described above by providing the entire non-human donor antibody heavy chain and light chain variable regions, including framework regions, in association with human immunoglobulin constant regions for both chains. It is anticipated that chimeric antibodies which retain additional non-human sequence relative to humanized antibodies of this invention may elicit a significant immune response in humans.

Such antibodies are useful in the prevention and treatment of IL4 mediated allergic disorders, as discussed below.

VI. Production of Fusion Proteins and Engineered Antibodies

Preferably, the variable light and/or heavy chain sequences and the CDRs of MAb 3B9 [SEQ ID NO: 16, 18, 20, 22, 24 and 26] or other suitable donor MABs (e.g., 6A1), and their encoding nucleic acid sequences, are utilized in the
5 construction of fusion proteins and engineered antibodies, preferably humanized antibodies, of this invention, by the following process. The same or similar techniques may also be employed to generate other embodiments of this invention.

A hybridoma producing a selected donor MAB, e.g., the murine antibody 3B9, is conventionally cloned, and the DNA of its heavy and light chain variable
10 regions obtained by techniques known to one of skill in the art, e.g., the techniques described in Sambrook *et al.*, Molecular Cloning (A Laboratory Manual), 2nd edition, Cold Spring Harbor Laboratory (1989). The variable heavy and light regions of 3B9 containing at least the CDRs and those portions of the acceptor MAB light and/or heavy variable domain framework region required in order to retain
15 donor MAB binding specificity, as well as the remaining immunoglobulin-derived parts of the antibody chain derived from a human immunoglobulin are obtained using polynucleotide primers and reverse transcriptase. The CDRs are identified using a known database and by comparison to other antibodies.

A mouse/human chimeric antibody may then be prepared and assayed for
20 binding ability. Such a chimeric antibody contains the entire non-human donor antibody V_H and V_L regions, in association with human Ig constant regions for both chains.

Homologous framework regions of a heavy chain variable region from a human antibody were identified using computerized databases, e.g., KABAT®, and
25 a human antibody having homology to 3B9 was selected as the acceptor antibody. The sequences of synthetic heavy chain variable regions containing the 3B9 CDRs within the human antibody frameworks were designed with optional nucleotide replacements in the framework regions to incorporate restriction sites. This designed sequence is then synthesized by overlapping oligonucleotides, amplified by
30 polymerase chain reaction (PCR), and corrected for errors.

A suitable light chain variable framework region was designed in a similar manner.

A humanized antibody may be derived from the chimeric antibody, or preferably, made synthetically by inserting the donor MAB CDRs from the heavy
35 and light chains appropriately within the selected heavy and light chain framework. Alternatively, a humanized antibody of the invention may be prepared using standard mutagenesis techniques. Thus, the resulting humanized antibody contains

human framework regions and donor MAb CDRs. There may be subsequent manipulation of framework residues. The resulting humanized antibody can be expressed in recombinant host cells, e.g., COS or CHO cells. Additional details of this procedure are provided in Example 4. Other humanized antibodies may be prepared using this technique on other suitable IL4-specific, neutralizing, high titer, non-human antibodies.

A conventional expression vector or recombinant plasmid is produced by placing these coding sequences for the fusion protein in operative association with conventional regulatory control sequences capable of controlling the replication and expression in, and/or secretion from, a host cell. Regulatory sequences include promoter sequences, e.g., CMV promoter, and signal sequences, which can be derived from other known antibodies. Similarly, a second expression vector is produced having a DNA sequence which encodes a complementary antibody light or heavy chain. Preferably this second expression vector is identical to the first except insofar as the coding sequences and selectable markers are concerned so to ensure as far as possible that each polypeptide chain is functionally expressed.

A selected host cell is co-transfected by conventional techniques with both the first and second vectors or simply transfected by a single vector to create the transfected host cell of the invention comprising both the recombinant or synthetic light and heavy chains. The transfected cell is then cultured by conventional techniques to produce the engineered antibody of the invention. The humanized antibody which includes the association of both the recombinant heavy chain and/or light chain is screened from culture by appropriate assay, such as ELISA or RIA. Similar conventional techniques may be employed to construct other fusion proteins and molecules of this invention.

Suitable vectors for the cloning and subcloning steps employed in the methods and construction of the compositions of this invention may be selected by one of skill in the art. For example, the conventional pUC series of cloning vectors, may be used. One vector used is pUC19, which is commercially available from supply houses, such as Amersham (Buckinghamshire, United Kingdom) or Pharmacia (Uppsala, Sweden). Additionally, any vector which is capable of replicating readily, has an abundance of cloning sites and marker genes, and is easily manipulated may be used for cloning. Thus, the selection of the cloning vector is not a limiting factor in this invention.

Similarly, the vectors employed for expression of the engineered antibodies according to this invention may be selected by one of skill in the art from any conventional vector. The vectors also contain selected regulatory sequences which

are in operative association with the DNA coding sequences of the immunoglobulin regions and capable of directing the replication and expression of heterologous DNA sequences in selected host cells, such as CMV promoters. These vectors contain the above described DNA sequences which code for the engineered antibody
5 or fusion molecule. Alternatively, the vectors may incorporate the selected immunoglobulin sequences modified by the insertion of desirable restriction sites for ready manipulation.

The expression vectors may also be characterized by marker genes suitable for amplifying expression of the heterologous DNA sequences, e.g., the mammalian
10 dihydrofolate reductase gene (DHFR) or neomycin resistance gene (neo^R). Other preferable vector sequences include a poly A signal sequence, such as from bovine growth hormone (BGH) and the betaglobin promoter sequence (betaglopro). The expression vectors useful herein may be synthesized by techniques well known to those skilled in this art.

15 The components of such vectors, e.g. replicons, selection genes, enhancers, promoters, signal sequences and the like, may be obtained from natural sources or synthesized by known procedures for use in directing the expression and/or secretion of the product of the recombinant DNA in a selected host. Other appropriate expression vectors of which numerous types are known in the art for
20 mammalian, bacterial, insect, yeast, and fungal expression may also be selected for this purpose.

The present invention also encompasses a cell line transfected with a recombinant plasmid containing the coding sequences of the engineered antibodies or fusion molecules hereof. Host cells useful for the cloning and other
25 manipulations of these cloning vectors are also conventional. However, most desirably, cells from various strains of *E. coli* are used for replication of the cloning vectors and other steps in the construction of fusion proteins of this invention.

Suitable host cells or cell lines for the expression of the engineered antibody or fusion protein of the invention are preferably a eukaryotic cell such as CHO,
30 COS, a fibroblast cell (e.g. 3T3), and myeloid cells among others, and most preferably a mammalian cell, such as a CHO cell or a myeloid cell. Human cells may be used, thus enabling the molecule to be modified with human glycosylation patterns. Alternatively, other eukaryotic cell lines may be employed. The selection of suitable mammalian host cells and methods for transformation, culture,
35 amplification, screening and product production and purification are known in the art. See, e.g., Sambrook *et al.*, cited above.

Bacterial cells may prove useful as host cells suitable for the expression of the recombinant MAbs of the present invention. However, due to the tendency of proteins expressed in bacterial cells to be in an unfolded or improperly folded form or in a non-glycosylated form, any recombinant MAb produced in a bacterial cell
5 would have to be screened for retention of antigen binding ability. If the molecule expressed by the bacterial cell was produced in a properly folded form, that bacterial cell would be a desirable host. For example, various strains of *E. coli* used for expression are well-known as host cells in the field of biotechnology. Various strains of *B. subtilis*, *Streptomyces*, other bacilli and the like may also be employed
10 in this method.

Where desired, strains of yeast cells known to those skilled in the art are also available as host cells, as well as insect cells, e.g. *Drosophila* and *Lepidoptera* and viral expression systems. See, e.g. Miller *et al.*, Genetic Engineering, 8:277-298, Plenum Press (1986) and references cited therein.

15 The general methods by which the vectors of the invention may be constructed, transfection methods required to produce the host cells of the invention, and culture methods necessary to produce the fusion protein or engineered antibody of the invention from such host cell are all conventional techniques. Likewise, once produced, the fusion proteins or engineered antibodies of the invention may be
20 purified from the cell culture contents according to standard procedures of the art, including ammonium sulfate precipitation, affinity columns, column chromatography, gel electrophoresis and the like. Such techniques are within the skill of the art and do not limit this invention.

Yet another method of expression of the humanized antibodies may utilize
25 expression in a transgenic animal, such as described in U. S. Patent No. 4,873,316. This relates to an expression system using the animal's casein promoter which when transgenically incorporated into a mammal permits the female to produce the desired recombinant protein in its milk.

Once expressed by the desired method, the engineered antibody is then
30 examined for *in vitro* activity by use of an appropriate assay. Presently conventional ELISA assay formats are employed to assess qualitative and quantitative binding of the engineered antibody to an IL4 epitope. Additionally, other *in vitro* assays, e.g. BIAcore [Pharmacia], may also be used to verify neutralizing efficacy prior to subsequent human clinical studies performed to
35 evaluate the persistence of the engineered antibody in the body despite the usual clearance mechanisms.

Following the procedures described for humanized antibodies prepared from 3B9, one of skill in the art may also construct humanized antibodies from other donor IL4 antibodies, variable region sequences and CDR peptides described herein. Engineered antibodies can be produced with variable region frameworks potentially
5 recognized as "self" by recipients of the engineered antibody. Minor modifications to the variable region frameworks can be implemented to effect large increases in antigen binding without appreciable increased immunogenicity for the recipient. Such engineered antibodies can effectively treat a human for IL4 mediated conditions. Such antibodies may also be useful in the diagnosis of such conditions.

10 VII. *Therapeutic/Prophylactic Uses*

This invention also relates to a method of treating humans experiencing an allergic disorder which comprises administering an effective dose of antibodies including one or more of the engineered antibodies or fusion proteins described herein, or fragments thereof.

15 The therapeutic response induced by the use of the molecules of this invention is produced by the binding to human IL4 and thus subsequently blocking IgE release. Thus, the molecules of the present invention, when in preparations and formulations appropriate for therapeutic use, are highly desirable for those persons experiencing an allergic response, such as an allergic rhinitis, conjunctivitis, atopic
20 dermatitis, atopic asthma, and anaphylactic shock.

The fusion proteins, antibodies, engineered antibodies or fragments thereof of this invention may also be used in conjunction with other antibodies, particularly human MAbs reactive with other markers (epitopes) responsible for the condition against which the engineered antibody of the invention is directed. Similarly MAbs
25 reactive with epitopes responsible for the condition in a selected animal against which the antibody of the invention is directed may also be employed in veterinary compositions.

The therapeutic agents of this invention are believed to be desirable for treatment of allergic conditions for from about 2 days to about 3 weeks, or as
30 needed. For example, longer treatments may be desirable when treating seasonal rhinitis or the like. This represents a considerable advance over the currently used infusion protocol with prior art treatments of IL4 mediated disorders. The dose and duration of treatment relates to the relative duration of the molecules of the present invention in the human circulation, and can be adjusted by one of skill in the art
35 depending upon the condition being treated and the general health of the patient.

The mode of administration of the therapeutic agent of the invention may be any suitable route which delivers the agent to the host. The fusion proteins,

antibodies, engineered antibodies, and fragments thereof, and pharmaceutical compositions of the invention are particularly useful for parenteral administration, i.e., subcutaneously, intramuscularly, intravenously, or intranasally.

Therapeutic agents of the invention may be prepared as pharmaceutical compositions containing an effective amount of the engineered (e.g., humanized) antibody of the invention as an active ingredient in a pharmaceutically acceptable carrier. In the prophylactic agent of the invention, an aqueous suspension or solution containing the engineered antibody, preferably buffered at physiological pH, in a form ready for injection is preferred. The compositions for parenteral administration will commonly comprise a solution of the engineered antibody of the invention or a cocktail thereof dissolved in an pharmaceutically acceptable carrier, preferably an aqueous carrier. A variety of aqueous carriers may be employed, e.g., 0.4% saline, 0.3% glycine, and the like. These solutions are sterile and generally free of particulate matter. These solutions may be sterilized by conventional, well known sterilization techniques (e.g., filtration). The compositions may contain pharmaceutically acceptable auxiliary substances as required to approximate physiological conditions such as pH adjusting and buffering agents, etc. The concentration of the antibody of the invention in such pharmaceutical formulation can vary widely, i.e., from less than about 0.5%, usually at or at least about 1% to as much as 15 or 20% by weight and will be selected primarily based on fluid volumes, viscosities, etc., according to the particular mode of administration selected.

Thus, a pharmaceutical composition of the invention for intramuscular injection could be prepared to contain 1 mL sterile buffered water, and between about 1 ng to about 100 mg, e.g. about 50 ng to about 30 mg or more preferably, about 5 mg to about 25 mg, of an engineered antibody of the invention. Similarly, a pharmaceutical composition of the invention for intravenous infusion could be made up to contain about 250 ml of sterile Ringer's solution, and about 1 to about 30 and preferably 5 mg to about 25 mg of an engineered antibody of the invention. Actual methods for preparing parenterally administrable compositions are well known or will be apparent to those skilled in the art and are described in more detail in, for example, Remington's Pharmaceutical Science, 15th ed., Mack Publishing Company, Easton, Pennsylvania.

It is preferred that the therapeutic agent of the invention, when in a pharmaceutical preparation, be present in unit dose forms. The appropriate therapeutically effective dose can be determined readily by those of skill in the art. To effectively treat an inflammatory disorder in a human or other animal, one dose of approximately 0.1 mg to approximately 20 mg per 70 kg body weight of a protein

or an antibody of this invention should be administered parenterally, preferably i.m. (intramuscularly). Such dose may, if necessary, be repeated at appropriate time intervals selected as appropriate by a physician during the inflammatory response.

5 The invention also encompasses the administration of the IL4 fusion proteins of this invention concurrently or sequentially with other antibodies or fusion proteins characterized by anti-IL4 activity, such as anti-tumor necrosis factor activity or other pharmaceutical activities compatible with the IL4 receptor binding ability of the fusion proteins of this invention. Such other antibodies are available commercially or can be designed in a manner similar to that described herein.

10 The fusion proteins and engineered antibodies of this invention may also be used in diagnostic regimens, such as for the determination of IL4 mediated disorders or tracking progress of treatment of such disorders. As diagnostic reagents, these fusion proteins may be conventionally labelled for use in ELISA's and other conventional assay formats for the measurement of IL4 levels in serum, plasma or
15 other appropriate tissue. The nature of the assay in which the fusion proteins are used are conventional and do not limit this disclosure.

The antibodies, engineered antibodies or fragments thereof described herein can be lyophilized for storage and reconstituted in a suitable carrier prior to use. This technique has been shown to be effective with conventional immunoglobulins and art-known lyophilization and reconstitution techniques can be employed.
20

The following examples illustrate various aspects of this invention including the construction of exemplary engineered antibodies and expression thereof in suitable vectors and host cells, and are not to be construed as limiting the scope of this invention. All amino acids are identified by conventional three letter or single
25 letter codes. All necessary restriction enzymes, plasmids, and other reagents and materials were obtained from commercial sources unless otherwise indicated. All general cloning ligation and other recombinant DNA methodology were as performed in T. Maniatis *et al.*, cited above, or the second edition thereof (1989), eds. Sambrook *et al.*, by the same publisher ("Sambrook *et al.*").

30

Example 1 - Production of MAb 3B9

A. Immunization procedure

Four mice (F1 hybrids of Balb/c and C57BL/6) were immunized subcutaneously with 50 µg recombinant *E. coli* human IL4 in Freund's complete adjuvant and 4 weeks later boosted intraperitoneally (i.p.) with 50 µg IL4 in Freund's incomplete adjuvant. On the basis of a good serum antibody titre to IL4 one mouse received further immunizations of 200 µg IL4 (i.p. in saline) at 8 weeks, two days later with 100 µg IL4 (i.p. in saline) and two days later with 50 µg IL4 (i.p. in saline). Two days following the final immunization a splenectomy was performed.

B. Fusion Procedure and Screening System

Mouse spleen cells were used to prepare hybridomas (by standard procedures, e.g. as described by Kohler et al, *Nature*, 256:495 (1975)) from which >250 clones of cells were screened for secretion of antibody to IL4, using the commercially available BIAcore system, and ELISA assays as described below, for IL4 binding. Five wells gave a positive response. Only 1 clone from mice, 3B9, was strongly positive. All secondary clones derived from 3B9 were positive.

Example 2 - ELISA Assays and Affinity Constants

A. ELISA

The screening assay, performed as follows, was designed to measure affinity for native human IL4. For experiment 1 aldehyde activated 96-well plates were coated with IL4 at 1 µg/mL, 100 µL/well in 0.1 M borate buffer, pH 8.5, and incubated overnight at RT. The hIL4 was covalently attached to the plate. IL4 solution was aspirated and non-specific binding (NSB) sites were blocked with 1% bovine serum albumin (BSA) in TBS buffer (50 mM Tris, 150 mM NaCl, 1 mM MgCl₂, 0.02% NaN₃, pH 7.4) for 60 minutes at 37°C. Following this and each of the following steps, the plate was washed 4 times in wash buffer (10 mM Tris, 150 mM NaCl, 0.05% Tween 20, 0.02% NaN₃, pH 7.4). Following this, 50 µL hybridoma medium (or purified 3B9 or Fab fragments) and 50 µL assay buffer (0.5% bovine gamma globulin in TBS buffer) was added and the plates were incubated for 60 minutes at 37°C. One hundred µL of biotinylated anti-mouse antibody was added per well in assay buffer and incubated as above. One hundred µL of alkaline phosphatase conjugated streptavidin was added per well and incubated (30 minutes at 37°C). One hundred µL/well PNP substrate was added and incubated 30 minutes at 37°C. Readings were taken at an optical density of 405 nm.

For experiment 2, streptavidin-coated plates (100 μ L/well, 1 μ g/mL in phosphate buffered saline (PBS)) were incubated overnight at 4°C and were assayed as follows. Streptavidin solution was aspirated, NSB sites blocked with 1% BSA in TBS buffer (60 minutes at 37°C). Following this step, and each of the steps which
 5 follow, the plates were washed four times in wash buffer. Fifty μ L biotinylated IL4 was added with 50 μ L assay buffer and incubated for 30 minutes at 37°C. Following this, 50 μ L purified 3B9 IgG or Fab fragment (or hybridoma medium) plus 50 μ L assay buffer was added, incubated 60 minutes at 37°C. One hundred μ L anti-mouse IgG alkaline phosphatase conjugate was added and incubated for 60
 10 minutes at 37°C. One hundred μ L PNP substrate was added and incubated 30 minutes at 37°C. The readings were taken as above.

B. Calculation of 3B9 Affinity for IL-4

Using the results of the experiments described above, and summarized as follows, the K_d for 3B9 was calculated as described in Beatty et al, J. Immunol. Methods, 100:173-179 (1987):
 15

$$K_{aff} = \frac{1}{2([Ab^*] - [Ab])}$$

Ab^* = concentration of Ab bound at 150 ng/ml biotinylated hIL4

Ab = concentration of Ab bound at 300 ng/ml biotinylated hIL4

20 Dissociation constants, K_d , were calculated from the relationship:

$$K_d = \frac{1}{K_{aff}}$$

Experiment 1: ELISA assay on a streptavidin coated 96-well plate (100 ng/well). $K_d = 2.2 \times 10^{-10}$ M (3B9 Fab)

25 Experiment 2: ELISA assay on a streptavidin coated 96-well plate (100 ng/well). $K_d = 1.4 \times 10^{-10}$ M (3B9 IgG)

C. Specificity

MAB 3B9 recognizes human IL4, but does not recognize bovine or murine IL4. One way to determine this is as follows. An ELISA can be performed using a
 30 96 well plate coated with anti-mouse IgG, and subsequently blocked with bovine serum albumin, upon which 50 μ L 3B9 (100 ng/mL), 25 μ L of non-human IL4, and 25 μ L biotin-IL4 were incubated for 60 minutes at 37°C, followed by a wash, streptavidin conjugated alkaline phosphatase and PNP.

Similarly, MAB 6A1 was found not to recognize bovine or murine IL4.

35

Example 3 - Humanized Antibody

One humanized antibody was designed to contain murine CDRs within a human antibody framework. This humanized version of the IL4 specific mouse antibody 3B9, was prepared by performing the following manipulations.

5

A. cDNA Cloning

cDNA clones were made of the 3B9 heavy and light chains from mRNA extracted out of the 3B9 hybridoma cell line [Example 1] using a Boehringer-Mannheim kit. Primers specific for either the mouse hinge region or kappa constant region were used for first strand synthesis.

10

The kappa chain primer is [SEQ ID NO: 29]:

5'-CTAACACTCATTCCTGTTGAAGCTCTTGACAATGGG-3'

The gamma heavy chain primer is [SEQ ID NO: 30]:

5'GTACATATGCAAGGCTTACAACCACAATC 3'.

15

The double stranded cDNA was cloned directly into plasmids pGEM7f+ [Promega] that were then transformed into *E. coli* DH5- α [Bethesda Research Labs].

B. DNA Sequencing

Eight heavy and one light chain murine cDNA clones from Part A above were sequenced. The results of sequencing of the variable regions of these clones are shown in SEQ ID NO: 1 and 2 and 3 and 4. Each clone contained amino acids known to be conserved among mouse heavy chain variable regions or light chain variable regions, and murine signal sequences. The CDR amino acid sequences are listed below.

20

The CDR regions for the heavy chain are SEQ ID NO: 22, 24 and 26, (amino acids 50-56, 71-86 and 119-129 of SEQ ID NO: 4). See Fig. 2. These sequences are encoded by SEQ ID NO: 21, SEQ ID NO: 23, and SEQ ID NO: 25, respectively. The CDR regions for the light chain are SEQ ID NO: 16, 18 and 20 (amino acids 45-58, 74-80, and 113-121 of SEQ ID NO: 2). See Fig. 1. These sequences are encoded by SEQ ID NO: 15, 17, and 19, respectively.

25

30

C. Selection of Human Frameworks

Following the cloning of 3B9, the amino acid sequences of the variable region (amino acids 21-132 of SEQ ID NO: 2 and amino acids 20 to 140 of SEQ ID NO: 4) were compared with the human immunoglobulin sequence database using the KABAT® and the SWISS databases in order to identify a human framework for both the heavy and light chains which would most closely match the murine parent in sequence homology. In addition to these searches for sequence homology, the

35

heavy and light chains were also evaluated against a positional database generated from structural models of the Fab domain to assess potential conflicts due to amino acid substitutions which might influence CDR presentation. For the present case, no obvious conflicts were detected in the structural search; hence, the DNA coding
5 deduced from the amino acid sequence homology searches was used.

The heavy chain framework regions of an antibody obtained from a human myeloma immunoglobulin (COR) was used [E. M. Press and N. M. Hogg, Biochem. J., 117:641-660 (1970)]. This sequence was found to be approximately 77% homologous (69.4% identity) to the 3B9 variable chain region at the amino acid
10 level.

For a suitable light chain variable framework region, the light chain variable framework sequence of the human antibody identified in H. G. Klobeck *et al.*, Nucl. Acids Res., 13:6515-6529 (1985) was used. The human antibody sequence was found to be approximately 80.2% homologous (72.0% identity) to the murine 3B9
15 variable light chain region at the amino acid level.

Given the murine 3B9 CDRs [SEQ ID NO: 15-26] and the sequence of the human antibody, a synthetic heavy chain was made and PCR performed to fill in and amplify the DNA. These sequences were synthesized by the following overlapping oligonucleotides and amplified by PCR. SEQ ID NO: 31-37 provides
20 five overlapping oligos and 2 PCR primers. Oligo 1 [SEQ ID NO: 31] is found spanning bases 5-121. Oligo 2 [SEQ ID NO: 32] is found spanning bases 122-241, and oligo 3 [SEQ ID NO: 33] is found spanning bases 242-361. The two bottom strand primers SEQ ID NO: 34 and SEQ ID NO: 35 span bases 134-110 and bases 253-230. Any errors in the mapped sequence which were inserted by PCR were
25 corrected. PCR was again performed using as the 5' primer nucleotides 1-25 SEQ ID NO: 36 and as the 3' primer nucleotides 361-341 SEQ ID NO: 37.

The synthetic variable region was ligated into the expression vector pCD along with the synthetic signal sequence SEQ ID NO: 5 and 6 from the chimeric heavy chain construction along with an IgG₁ human constant region. The synthetic
30 V_H and signal sequence nucleotide and amino acid sequences are provided in Fig. 4 [SEQ ID NOS: 11 and 12]. The amino acid sequences of the CDRs [SEQ ID NOS: 22, 24 and 26] are identical to the murine 3B9 CDRs. However, the coding sequences for these CDRs [SEQ ID NOS: 54, 55 and 56] differ from the murine 3B9 coding sequences [SEQ ID NOS: 21, 23 and 25]. The resulting expression
35 vector, IL4hzhcl-1-Pcd is shown in Fig. 9.

The CDR gene regions of a pre-existing light chain framework were restriction digest removed and replaced with the following synthetic IL-4 CDR genes, which were synthetically made.

For CDR1:

- 5 SEQ ID NO: 38: 5'CTAGCTGTGTCTCTGGGCGAGAGGGCCACCATCAAC
TGCAAGG 3'
SEQ ID NO: 39: CCTTGCAGTTGATGGTGGCCCTCTCGCCCAGAGACACAG
SEQ ID NO: 40: TCGAGAGGCCTCCCAAAGTGTTGATTATGATGGTGATAG
TTATATGAACTGGTATCAGCAGAAACCC
10 SEQ ID NO: 41:
GGGTTTCTGCTGATACCAGTTCATATAACTATCACCATCATA
ATCAACACTTTGGGAGGCCTC

For CDR2:

- SEQ ID NO: 44:
15 GGGCAGCCTCCTAAGTTGCTCATTTACGCTGCATCCAATCTA
GAATCTGGGGTAC
SEQ ID NO: 45:
CCCAGATTCTAGATTGGATGCAGCGTAAATGAGCAACTTAGG
AGGCTGCCC

20 For CDR3:

- SEQ ID NO: 42:
ATACTACTGTCAGCAAAGTAATGAGGATCCTCCGAGGTTCCGG
CGGAGGGAC
SEQ ID NO: 43:
25 CTTGGTCCCTCCGCCGAACCTCGGAGGATCCTCATTACTTTG
CTGACAGTAGT

- The synthetic V_L and signal sequence nucleotide and amino acid sequences are provided in Fig. 5 [SEQ ID NOS: 13 and 14]. The amino acid sequences of the first two CDRs [SEQ ID NOS: 16 and 18] are identical to the corresponding murine 3B9 CDRs. However, the coding sequence for the first CDR [SEQ ID NO: 53] differs from the murine 3B9 coding sequence [SEQ ID NO: 15]. Further, in the last CDR, two humanized constructs of the 3B9 amino acid sequence were constructed. One, [SEQ ID NO: 28], differs by a single amino acid [SEQ ID NO: 20] from the native murine 3B9 sequence. SEQ ID NO: 28 is encoded by SEQ ID NO: 27. The synthetic variable light regions were ligated into the expression vector along with the signal sequence [SEQ ID NOS: 7 and 8]. One of the resulting expression vectors, IL4hzlc1-O-Pcn is illustrated in Fig. 10.

These synthetic variable light and/or heavy chain sequences are employed in the construction of a humanized antibody.

Example 4 - Expression of Humanized MAb in COS and CHO cells

5 pUC18 subclones for the V_H were made to add a signal sequence originally obtained from a human antibody SEQ ID NO: 5. For the V_L, pUC18 subclones were made to add a signal sequence SEQ ID NO: 7.

The humanized heavy chain, derived from an IgG₁ isotype, exhibits 89.3% homology (84.3% identity) at the amino acid level with the murine heavy chain from 3B9. This synthetic V_H is provided in amino acids 20-141 of SEQ ID NOS: 11 and 12.

The humanized light chain, a human kappa chain, shows 92.0% homology (86.6% identity) with 3B9 at the amino acid level. This synthetic V_L [amino acids 21 to 131 of SEQ ID NOS: 13 and 14] containing the 3B9 CDRs was designed and synthesized as described above for the synthetic heavy chains.

The DNA fragments containing their respective signal linked to either the humanized heavy or light variable regions were inserted into pUC19-based mammalian cell expression plasmids containing CMV promoters and the human heavy chain or human light chain constant regions of the chimera produced in Example 5 below, by conventional methods [Maniatis *et al.*, cited above] to yield the plasmids IL4hzhc1-1Pcd (heavy chain) [Figure 9] and IL4hzlc1-o-Pcn (light chain) [Figure 10]. The HZHC and HZLC plasmids are co-transfected into COS cells and supernatants assayed by the ELISA described immediately above for the presence of humanized antibody after three and five days. Another humanized antibody was constructed but with an IgG4 isotype.

The above example describes the preparation of an exemplary engineered antibody. Similar procedures may be followed for the development of other engineered antibodies, using other anti-IL4 antibodies (e.g., 6A1 - Example 7) developed by conventional means.

Example 5 - Construction of Chimeric Antibody

A. A chimeric heavy chain was constructed by isolating the murine variable heavy chain region from the original mouse MAb 3B9 as an EcoRI-BstEII restriction fragment. A small DNA oligomer was designed and synthesized to link the mouse variable region with the human IgG1 constant region (BstEII - ApaI):
 35 5' primer: SEQ ID NO: 50: GTCACCGTCTCCTCAGCTAGCACCAAGGGGC
 3' primer: SEQ ID NO: 51: CTTGGTGCTAGCTGAGGAGACG

These two fragments were ligated into plasmid pCD (See Fig. 7)(digested with EcoRI and ApaI) that already encodes the human IgG1 constant region. This clone did not express; therefore, the wild-type 5'UTR and signal sequence were deleted and replaced with SEQ ID NO:5 and 6.

5 Because a convenient restriction endonuclease site was not available at the 3' end of the signal sequence, a BstEII site was introduced (i.e., a silent mutation) via PCR. The following PCR primers were used:

SEQ ID NO: 48: 5' primer: 5'CAGGTTACCCCTGAAAGAGTC 3'

SEQ ID NO: 49: 3' primer: 5'GAAGTAGTCCTTGACCAG 3'

10 A BstEII - PstI restriction fragment was then isolated from this plasmid. A new signal sequence and 5'UTR were then designed and synthesized having EcoRI and BstEII ends.

SEQ ID NO: 46: 5' primer: AATTCGAGGACGCCAGCAACATGGTGTGCA
GACCCAGGTCTTCATTTCTCTGTTGCTCTGGATCTCTGGTGCCTACGGGC

15 AG

SEQ ID NO: 47: 3' primer: GTAACCTGCCCGTAGGCACCAGAGATCCAGA
GCAACAGAGAAATGAAGACCTGGGTCTGCAACACCATGTTGCTGGCGTC
CTCG

The chimeric light chain was constructed by applying the PCR
20 technique to the original murine 3B9 light chain that was cloned into pGEM72f(+) [Promega]. The primers utilized were the commercially available pUC18 universal reverse primer at the 5' end (EcoRI) and a 3' primer that introduces a NarI site [5'CATCTAGATGGCG CCGCCACAGTACGTTTGATCTCCAGCTTGGTCCC3' SEQ ID NO: 52], used to fuse the mouse variable region to the human constant
25 region. This variable region was then ligated into the expression vector pCDN (EcoRI NarI) (Fig. 8) that already contains the human kappa region.

Media supernatants were collected three and five days later and assayed by the ELISA described as follows: ELISA plates were coated with 0.1µg of a goat antibody specific for the Fc region of human antibodies. The media
30 supernatants were added for one hour. A horseradish peroxidase conjugated goat antibody specific for an entire human IgG antibody was added. This was followed by addition of ABTS peroxidase substrate (Kirkegaard & Perry Laboratories Inc., Gaithersburg, MD) for one hour. Expression of the chimeric antibody was detected. In a second ELISA the COS cell supernatants containing the chimeric antibody
35 bound specifically to recombinant human IL4 protein. This result confirmed that genes coding for an antibody specific for IL4 had been cloned.

B. A humanized heavy chain can also be obtained from this chimeric heavy chain. The humanized heavy chain was designed from by inserting the murine CDRs into a human framework. The chosen human framework was as described above, the most homologous protein sequence in the Swiss protein data
5 based to the murine 3B9 V_H (amino acids 20-140 of SEQ ID NO: 4). This humanized heavy chain sequence (EcoRI ApaI) was made synthetically and PCR performed to fill in and amplify DNA as described above. This synthetic variable region was ligated into the the expression vector pCD (EcoRI ApaI) together with the synthetic signal sequence SEQ ID NOS: 5 and 6 from the chimeric heavy chain
10 construction and an IgG₁ human constant region.

Similarly, a humanized light chain can be derived from the chimeric light chain as described for the heavy chain. This gene (EcoRV NarI) was also made synthetically. The humanized V_L was ligated into the expression vector pCN, digested with EcoRI NarI, along with a signal sequence (EcoRI EcoRV). The
15 expression vector provided the human kappa constant region.

Example 6 - Purification and Thermodynamics - Humanized MAb

Purification of CHO expressed chimeric and humanized 3B9 can be achieved by conventional protein A (or G) affinity chromatography followed by ion
20 exchange and molecular sieve chromatography. Similar processes have been successfully employed for the purification to >95% purity of other MAbs (e.g., to respiratory syncytial virus and malaria circumsporozoite antigens).

The affinity and detailed thermodynamics of IL4 binding to humanized MAb 3B9 and murine 3B9 (Example 1) were determined by titration microcalorimetry.
25 This method measures binding reactions by virtue of their intrinsic heats of reaction (see, e.g., Wiseman et al., Anal. Biochem., 179:131-137 (1989)). The affinity of both MAbs was found to be too tight to measure directly at ambient temperature. Thus, a thermodynamic approach was taken: i) the affinity was measured at 60°C, where it is weak enough to be measured directly; and (ii) the temperature-dependence of the
30 binding enthalpy was measured from 30-60°C. Together, these data allow calculation of the affinity over a wide range of temperatures using the Gibbs-Helmholz equation.

A summary of the IL4 binding thermodynamics of the humanized and murine 3B9 antibodies are presented in Table 1. Based upon the changes in free
35 energy, enthalpy, entropy and heat capacity of the two MAbs, the binding thermodynamics are indistinguishable.

Table 1

Thermodynamics of hIL-4 binding to Humanized 3B9 and Murine 3B9 at pH 7.4, 150 mM NaCl, and 25°C.

mAb	K _d	ΔG	ΔH	-TΔS	ΔC
	picomolar	kcal/ mol IL4	kcal/ mol IL4	kcal/ mol IL4	cal/mol IL4/°K
humanized 3B9	11	-13.6±0.6	-21.8±2	8.2±2.1	-580±160
murine 3B9	19	-13.3±0.6	-20.5±1	7.2±1.2	-660±200

IL-4 affinities of humanized 3B9 and murine 3B9 were measured in quadruplicate and duplicate, respectively.

Example 7 - Production and Characterization of Rat MAb

MAb 6A1, chosen for high affinity binding, was derived from an immunized rat, using the same immunization protocol as described for the mouse in Example 1.

6A1 was selected from hybridomas (specifically, hybridoma 3426A11C1B9) prepared from rats immunized with human IL4.

The K_d for 6A1 was calculated as described in Beatty *et al*, *J. Immunol. Methods*, 100:173-179 (1987) to be 2×10^{-10} M.

Hybridoma 3426A11C1B9 was deposited October 6, 1993 with the European Collection of Animal Cell Cultures (ECACC), Public Health Laboratory Service Centre for Applied Microbiology & Research, Porton Down, Salisbury, Wiltshire, SP4 0JG, United Kingdom, under accession number 93100620, and has been accepted as a patent deposit, in accordance with the Budapest Treaty of 1977 governing the deposit of microorganisms for the purposes of patent procedure.

Example 8 - Biological Activity of MAbs: 3B9 (humanized), 3B9 (Murine) and 6A1

The following assays were performed using the procedures described below.

A. Binding to Glycosylated rhIL4

The above-identified antibodies were raised to non-glycosylated recombinant human IL4 (rhIL4) which was produced in *E. coli*. Because native human IL4 is glycosylated, it was important to confirm binding to material secreted by a

mammalian cell line. 3B9 binds equally well to both glycosylated and non-glycosylated human recombinant IL4, and is not therefore directed to an epitope that would be masked on natural human IL4.

B. Inhibition of IL4 Binding to Receptor

- 5 The ability of 3B9 to inhibit the binding of IL4 to its receptor was studied using ^{125}I -rhIL4 binding to the gibbon cell line, MLA [ATCC TIB201], that bears approximately 6000 receptors per cell. MLA cells were incubated with ^{125}I -IL4 for 30 minutes at 37°C. Uptake of radioactivity was determined in a gamma counter after separation of cell bound ^{125}I -IL4 by centrifugation through an oil-gradient.
- 10 Non-specific binding was determined by incubating in the presence of a 100-fold molar excess of unlabelled IL4 [Park et al, J. Exp. Med., 166:476-488 (1987)]. The IC_{50} value for unlabeled IL4 in this assay was 22 pM when the amount of (added) IL4 was 83 pM. For intact murine (IgG) 3B9 the IC_{50} was 63 pM, and 93 pM for the Fab fragment. At another concentration of IL4 (218 pM), the assay amount for
- 15 murine (IgG) 3B9 was 109 pM.

C. Inhibition of Lymphocyte Proliferation

- Using the method described in Spits et al, J. Immunol., 139:1142-1147 (1987), human peripheral blood lymphocytes are incubated for three days with phytohemagglutinin, a T cell mitogen, to upregulate the IL4 receptor. The resultant
- 20 blast cells are then stimulated for three days further with IL4. Proliferation is measured by the incorporation of ^3H thymidine. B cell proliferation was measured by the assay of Callard et al, in Lymphokines and Interferons. A Practical Approach, Ch. 19, p. 345, modified as follows. Purified human tonsillar B cells are stimulated for 3 days with IL4 and immobilized anti-IgM. Proliferation is measured
- 25 by the incorporation of ^3H thymidine.

- 3B9 (murine) inhibited ^3H -thymidine incorporation by human peripheral blood T lymphocytes stimulated with 133 pM IL4 and human tonsillar B lymphocytes stimulated by 167 pM IL4. IL2-stimulated T lymphocytes were not affected. The IC_{50} for inhibition of T cell proliferation was 30 pM, and for B cell
- 30 proliferation 103 pM. The corresponding values for the Fab fragment of 3B9 (murine) were 108 and 393 pM.

D. Inhibition of CD23 Induction

- CD23 is the low affinity receptor for IgE (FcERII) and is induced on the membrane of resting B lymphocytes by low concentrations of IL4 as a necessary
- 35 prerequisite for IgE production. Purified human tonsillar B cells are stimulated for 2 days with IL4. The percentage of cells expressing the CD23 receptor are determined by flow cytometry [Defrance et al, J. Exp. Med., 165:1459-1467

(1987)]. 3B9 (murine) inhibited CD23 expression on human tonsil B lymphocytes stimulated with 8.3 pM IL4 with an IC_{50} value of 136 pM.

E. Inhibition of IgE Secretion

Unlike other assays where IL4 was added at EC_{50} concentrations [Pere et al, 5 Proc. Natl. Acad. Sci., 85:6880-6884 (1988)], IgE secretion was investigated in the presence of concentrations of IL4 giving maximal secretion in order to reduce the variability inherent in this system. T cell proliferation was measured as follows. Human peripheral blood lymphocytes are incubated with IL4 for between 10-18, preferably 12, days. The concentration of IgE in the culture supernatant is 10 determined by ELISA.

IgE secretion was inhibited by 3B9 (murine), and the Fab fragment of 3B9, in the presence of 1.7 nM IL4 giving IC_{50} values of 1.9 and 5.0 nM respectively. The experiment was repeated using a lower concentration of IL4, 667 pM, which reduced the IC_{50} value to 0.65 nM for 3B9 (murine). The molar ratio of antibody 15 (IgG) to IL4 remained unchanged (1:1) over the concentration ranges examined.

F. Summary and Interpretation of Data

The molar ratios of IL4 to various MAbs required for 50% inhibition of function in bioassays is given in Table 2.

Table 2	Comparative activity of mAbs 3B9, 6A1 and Humanized 3B9 [IgG1 and IgG4 variants] in IL-4 dependent bioassays				
Assay	IC50 (pM) [range] n				
	Murine 3B9	Murine 3B9 (Fab)	Rat 6A1	Humanized 3B9	
				IgG1	IgG4*
RBA	63 [17-109] ₂	93	>50000		
T cell	30 [10-40] ₄	108	87	44[30-56] ₃	40
B cell	103 [79-120] ₃	393	187	47 [10-80] ₃	79
CD23 induction	136 [53-272] ₄	216	80	333	
IgE synthesis	658 [370-1070] ₆	1170	623 [412-833] ₂	54 [35-83] ₃	406

n = number of separate tests carried out.

*The IgG1 and IgG4 variants were assayed at different times.

In all assays, except IgE secretion, IL4 was added at approximate ED₅₀ concentrations. The molar ratios of antibody to IL-4 required for 50% inhibition were similar for humanized 3B9, murine 3B9, and 6A1 in the two lymphocyte proliferation assays, but higher for humanized 3B9 in the CD23 induction assay. The latter is a particularly sensitive assay apparently requiring very low (~ 5%) receptor occupancy (Kruse et al., EMBO J, 12:5121 1993) and, as is evident from the results obtained with murine 3B9, subject to inter assay variation.

A comparison of the activities of rat 6A1 and murine 3B9 demonstrated a similar profile of functional effects, but an unexpected failure of 6A1 to fully inhibit the binding of radioiodinated IL4 to its receptor. The radioiodinated IL4 used in the receptor binding assay is thought to be iodinated at the accessible tyrosine, residue 124. When the ability of 6A1 to inhibit CD23 expression induced by either

unlabelled or iodinated IL4 was compared, it was found that inhibition was less efficient against iodinated ligand. These results indicate that 6A1 binds to IL4 in the region of, but not specifically to, tyrosine 124.

Thus on current data, 6A1 is a neutralizing antibody of high affinity, binding to a very different region of IL4 than 3B9.

Example 9 - Pharmacokinetics

The pharmacokinetics of humanized 3B9 was investigated in the male Sprague Dawley rat. Humanized 3B9 was administered to four animals as an *iv* bolus dose at 1 mg/kg, blood sampling was continued for 5 weeks post dosing. Plasma humanized 3B9 concentrations were determined using an IL-4/anti-human IgG sandwich ELISA designed to confirm not only the presence of circulating human IgG but also its ability to bind to recombinant human IL-4.

Results from this study are summarized in Table 3.

Table 3
Pharmacokinetics of Humanized 3B9 in male Sprague-Dawley Rats
(dose: 1 mg/kg *iv* bolus)

	Clp (mL/h/kg)
Rat 1	0.442
Rat 2	0.655
Rat 3	0.555
Rat 4	0.447
Mean	0.525
SD	0.101

Abbreviation of the pharmacokinetic parameter is as follows: Clp, apparent plasma clearance.

Data indicated that inter-animal variability was relatively small and disappearance of humanized 3B9 from plasma appeared to be biphasic. The apparent plasma clearance was low (0.5 mL/h/kg). The half-life appeared to be 11 days. Thus, the pharmacokinetic characteristics of the CHO cell-derived humanized 3B9 are consistent with other humanized monoclonal antibodies in rats. The long circulating half life of humanized 3B9 in the rat also suggests that when

administered to man, humanized 3B9 is likely to be effective over an extended period of time.

5 Numerous modifications and variations of the present invention are included in the above-identified specification and are expected to be obvious to one of skill in the art. For example, human framework regions or modifications thereof, other than the exemplary antibodies described above, may be used in the construction of humanized antibodies. Such modifications and alterations to the compositions and processes of the present invention are believed to be encompassed in the scope of
10 the claims appended hereto.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT: Holmes, Stephen D.
Gross, Mitchell S.
Sylvester, Daniel R.

(ii) TITLE OF INVENTION: Recombinant IL4 Antibodies Useful in
Treatment of IL4 Mediated Disorders

(iii) NUMBER OF SEQUENCES: 58

(iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: SmithKline Beecham Corporation
(B) STREET: Corporate Intellectual Property, UW2220 - 709
Swedeland Rd.
(C) CITY: King of Prussia
(D) STATE: PA
(E) COUNTRY: USA
(F) ZIP: 19406-2799

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: PatentIn Release #1.0, Version #1.25

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER: US 08/117,366
(B) FILING DATE: 07-SEP-1993
(C) CLASSIFICATION:

(A) APPLICATION NUMBER: US 08/136,783
(B) FILING DATE: 14-OCT-1993
(C) CLASSIFICATION:

(viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: Sutton, Jeffrey A.

(C) REFERENCE/DOCKET NUMBER: P50186-2

(B) TELEFAX: (215) 270-5090

(D) TOPOLOGY: unknown

(B) LOCATION: 1..396

GTG TCT CTA GGG CAG AGG GCC ACC ATC TCC TGC AAG GCC AGC CAA AGT 144
Val Ser Leu Gly Gln Arg Ala Thr Ile Ser Cys Lys Ala Ser Gln Ser

35	40	45	
GTT GAT TAT GAT GGT GAT AGT TAT ATG AAC TGG TAC CAA CAG AAA CCA			192
Val Asp Tyr Asp Gly Asp Ser Tyr Met Asn Trp Tyr Gln Gln Lys Pro			
50	55	60	
GGA CAG CCA CCC AAA CTC CTC ATC TAT GCT GCA TCC AAT CTA GAA TCT			240
Gly Gln Pro Pro Lys Leu Leu Ile Tyr Ala Ala Ser Asn Leu Glu Ser			
65	70	75	80
GGG ATC CCA GCC AGG TTT AGT GGC AGT GGG TCT GGG ACA GAC TTC ACC			288
Gly Ile Pro Ala Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr			
85	90	95	
CTC AAC ATC CAT CCT GTG GAG GAG GAG GAT GCT GCA ACC TAT TAC TGT			336
Leu Asn Ile His Pro Val Glu Glu Glu Asp Ala Ala Thr Tyr Tyr Cys			
100	105	110	
CAG CAA AGT AAT GAG GAT CCT CCG ACG TTC GGT GGA GGC ACC AAG CTG			384
Gln Gln Ser Asn Glu Asp Pro Pro Thr Phe Gly Gly Gly Thr Lys Leu			
115	120	125	
GAA ATC AAA CGG			396
Glu Ile Lys Arg			
130			

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 132 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met	Glu	Thr	Asp	Thr	Ile	Leu	Leu	Trp	Val	Leu	Leu	Leu	Trp	Val	Pro
1					5				10				15		

Gly Ser Thr Gly Asp Ile Val Leu Thr Gln Ser Pro Ala Ser Leu Ala
 20 25 30
 Val Ser Leu Gly Gln Arg Ala Thr Ile Ser Cys Lys Ala Ser Gln Ser
 35 40 45
 Val Asp Tyr Asp Gly Asp Ser Tyr Met Asn Trp Tyr Gln Gln Lys Pro
 50 55 60
 Gly Gln Pro Pro Lys Leu Leu Ile Tyr Ala Ala Ser Asn Leu Glu Ser
 65 70 75 80
 Gly Ile Pro Ala Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr
 85 90 95
 Leu Asn Ile His Pro Val Glu Glu Glu Asp Ala Ala Thr Tyr Tyr Cys
 100 105 110
 Gln Gln Ser Asn Glu Asp Pro Pro Thr Phe Gly Gly Gly Thr Lys Leu
 115 120 125
 Glu Ile Lys Arg
 130

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 483 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 64..483

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

GAATTCGCGG CCGCTATGCA GGGACAATCA GCAGCAGCAA TGAGGAAGTA AGCCTGTGCA	60
GAT ATG AAC AGG CTT ACT TCC TCA TTG CTG CTG CTG ATT GTC CCT GCA	108
Met Asn Arg Leu Thr Ser Ser Leu Leu Leu Leu Ile Val Pro Ala	
1 5 10 15	
TAT GTC CTG TCC CAG GTT ACT CTG AAA GAG TCT GGC CCT GGG ATA TTG	156
Tyr Val Leu Ser Gln Val Thr Leu Lys Glu Ser Gly Pro Gly Ile Leu	
20 25 30	
CAG CCC TCC CAG ACC CTC AGT CTG ACT TGT TCT TTC TCT GGG TTT TCA	204
Gln Pro Ser Gln Thr Leu Ser Leu Thr Cys Ser Phe Ser Gly Phe Ser	
35 40 45	
CTG AGC ACT TCT GGT ATG GGT GTG AGC TGG ATT CGT CAG CCT TCA GGA	252
Leu Ser Thr Ser Gly Met Gly Val Ser Trp Ile Arg Gln Pro Ser Gly	
50 55 60	
AAG GGT CTG GAG TGG CTG GCA CAC ATT TAC TGG GAT GAT GAC AAG CGC	300
Lys Gly Leu Glu Trp Leu Ala His Ile Tyr Trp Asp Asp Asp Lys Arg	
65 70 75	
TAT AAC CCA TCC CTG AAG AGC CGG CTC ACA ATC TCC AAG GAT ACC TCC	348
Tyr Asn Pro Ser Leu Lys Ser Arg Leu Thr Ile Ser Lys Asp Thr Ser	
80 85 90 95	
AGC AAC CAG GTA TTC CTC AAG ATC ACC AGT GTG GAC ACT GCA GAT ACT	396
Ser Asn Gln Val Phe Leu Lys Ile Thr Ser Val Asp Thr Ala Asp Thr	
100 105 110	
GCC ACA TAC TAC TGT GCT CGA AGA GAG ACT GTG TTC TAC TGG TAC TTC	444
Ala Thr Tyr Tyr Cys Ala Arg Arg Glu Thr Val Phe Tyr Trp Tyr Phe	
115 120 125	
GAT GTC TGG GGC GCA GGG ACC ACG GTC ACC GTC TCC TCA	483
Asp Val Trp Gly Ala Gly Thr Thr Val Thr Val Ser Ser	

130

135

140

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 140 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Met Asn Arg Leu Thr Ser Ser Leu Leu Leu Leu Ile Val Pro Ala Tyr
 1 5 10 15
 Val Leu Ser Gln Val Thr Leu Lys Glu Ser Gly Pro Gly Ile Leu Gln
 20 25 30
 Pro Ser Gln Thr Leu Ser Leu Thr Cys Ser Phe Ser Gly Phe Ser Leu
 35 40 45
 Ser Thr Ser Gly Met Gly Val Ser Trp Ile Arg Gln Pro Ser Gly Lys
 50 55 60
 Gly Leu Glu Trp Leu Ala His Ile Tyr Trp Asp Asp Asp Lys Arg Tyr
 65 70 75 80
 Asn Pro Ser Leu Lys Ser Arg Leu Thr Ile Ser Lys Asp Thr Ser Ser
 85 90 95
 Asn Gln Val Phe Leu Lys Ile Thr Ser Val Asp Thr Ala Asp Thr Ala
 100 105 110
 Thr Tyr Tyr Cys Ala Arg Arg Glu Thr Val Phe Tyr Trp Tyr Phe Asp
 115 120 125
 Val Trp Gly Ala Gly Thr Thr Val Thr Val Ser Ser
 130 135 140

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 60 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 1..60

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

ATG GTG TTG CAG ACC CAG GTC TTC ATT TCT CTG TTG CTC TGG ATC TCT	48
Met Val Leu Gln Thr Gln Val Phe Ile Ser Leu Leu Leu Trp Ile Ser	
1 5 10 15	
GGT GCC TAC GGG	60
Gly Ala Tyr Gly	
20	

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

Met Val Leu Gln Thr Gln Val Phe Ile Ser Leu Leu Leu Trp Ile Ser

1 5 10 15
Gly Ala Tyr Gly
 20

(2) INFORMATION FOR SEQ ID NO:7:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 57 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

- (A) NAME/KEY: CDS
 (B) LOCATION: 1..57

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

ATG GGA TGG AGC TGT ATC ATC CTC TTC TTG GTA GCA ACA GCT ACA GGT	48
Met Gly Trp Ser Cys Ile Ile Leu Phe Leu Val Ala Thr Ala Thr Gly	
1 5 10 15	
GTC CAC TCC	57
Val His Ser	

(2) INFORMATION FOR SEQ ID NO:8:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 19 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

Met Gly Trp Ser Cys Ile Ile Leu Phe Leu Val Ala Thr Ala Thr Gly
 1 5 10 15

Val His Ser

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 423 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 1..423

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

ATG GTG TTG CAG ACC CAG GTC TTC ATT TCT CTG TTG CTC TGG ATC TCT 48
 Met Val Leu Gln Thr Gln Val Phe Ile Ser Leu Leu Trp Ile Ser
 1 5 10 15

GGT GCC TAC GGG CAG GTT ACC CTG AAA GAG TCT GGC CCT GGG ATA TTG 96
 Gly Ala Tyr Gly Gln Val Thr Leu Lys Glu Ser Gly Pro Gly Ile Leu
 20 25 30

CAG CCC TCC CAG ACC CTC AGT CTG ACT TGT TCT TTC TCT GGG TTT TCA 144
 Gln Pro Ser Gln Thr Leu Ser Leu Thr Cys Ser Phe Ser Gly Phe Ser
 35 40 45

CTG AGC ACT TCT GGT ATG GGT GTG AGC TGG ATT CGT CAG CCT TCA GGA 192
 Leu Ser Thr Ser Gly Met Gly Val Ser Trp Ile Arg Gln Pro Ser Gly
 50 55 60

GAT GTC TGG GGC GCA GGG ACC ACG GTC ACC GTC TCC TCA 423
Asp Val Trp Gly Ala Gly Thr Thr Val Thr Val Ser Ser
130 135 140

Gln Pro Ser Gln Thr Leu Ser Leu Thr Cys Ser Phe Ser Gly Phe Ser
 35 40 45

 Leu Ser Thr Ser Gly Met Gly Val Ser Trp Ile Arg Gln Pro Ser Gly
 50 55 60

 Lys Gly Leu Glu Trp Leu Ala His Ile Tyr Trp Asp Asp Asp Lys Arg
 65 70 75 80

 Tyr Asn Pro Ser Leu Lys Ser Arg Leu Thr Ile Ser Lys Asp Thr Ser
 85 90 95

 Ser Asn Gln Val Phe Leu Lys Ile Thr Ser Val Asp Thr Ala Asp Thr
 100 105 110

 Ala Thr Tyr Tyr Cys Ala Arg Arg Glu Thr Val Phe Tyr Trp Tyr Phe
 115 120 125

 Asp Val Trp Gly Ala Gly Thr Thr Val Thr Val Ser Ser
 130 135 140

(2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 423 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 1..423

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

ATG GTG TTG CAG ACC CAG GTC TTC ATT TCT CTG TTG CTC TGG ATC TCT

48

Met Val Leu Gln Thr Gln Val Phe Ile Ser Leu Leu Leu Trp Ile Ser	
1 5 10 15	
GGT GCC TAC GGG CAG GTT ACC CTG CGT GAA TCC GGT CCG GCA CTA GTT	96
Gly Ala Tyr Gly Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val	
20 25 30	
AAA CCG ACC CAG ACC CTG ACG TTA ACC TGC ACC TTC TCC GGT TTC TCC	144
Lys Pro Thr Gln Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser	
35 40 45	
CTG TCG ACC TCC GGT ATG GGT GTT TCC TGG ATC CGT CAG CCG CCG GGT	192
Leu Ser Thr Ser Gly Met Gly Val Ser Trp Ile Arg Gln Pro Pro Gly	
50 55 60	
AAA GGT CTA GAA TGG CTG GCT CAC ATC TAC TGG GAC GAC GAC AAA CGT	240
Lys Gly Leu Glu Trp Leu Ala His Ile Tyr Trp Asp Asp Asp Lys Arg	
65 70 75 80	
TAC AAC CCG AGC CTG AAA TCC CGT CTG ACG ATA TCC AAA GAC ACC TCC	288
Tyr Asn Pro Ser Leu Lys Ser Arg Leu Thr Ile Ser Lys Asp Thr Ser	
85 90 95	
CGT AAC CAG GTT GTT CTG ACC ATG ACT AAC ATG GAC CCG GTT GAC ACC	336
Arg Asn Gln Val Val Leu Thr Met Thr Asn Met Asp Pro Val Asp Thr	
100 105 110	
GCT ACC TAC TAC TGC GCT CGA CGC GAA ACC GTT TTC TAC TGG TAC TTC	384
Ala Thr Tyr Tyr Cys Ala Arg Arg Glu Thr Val Phe Tyr Trp Tyr Phe	
115 120 125	
GAC GTT TGG GGT CGT GGT ACC CCA GTT ACC GTG AGC TCA	423
Asp Val Trp Gly Arg Gly Thr Pro Val Thr Val Ser Ser	
130 135 140	

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 141 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

Met Val Leu Gln Thr Gln Val Phe Ile Ser Leu Leu Leu Trp Ile Ser
 1 5 10 15

Gly Ala Tyr Gly Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val
 20 25 30

Lys Pro Thr Gln Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser
 35 40 45

Leu Ser Thr Ser Gly Met Gly Val Ser Trp Ile Arg Gln Pro Pro Gly
 50 55 60

Lys Gly Leu Glu Trp Leu Ala His Ile Tyr Trp Asp Asp Asp Lys Arg
 65 70 75 80

Tyr Asn Pro Ser Leu Lys Ser Arg Leu Thr Ile Ser Lys Asp Thr Ser
 85 90 95

Arg Asn Gln Val Val Leu Thr Met Thr Asn Met Asp Pro Val Asp Thr
 100 105 110

Ala Thr Tyr Tyr Cys Ala Arg Arg Glu Thr Val Phe Tyr Trp Tyr Phe
 115 120 125

Asp Val Trp Gly Arg Gly Thr Pro Val Thr Val Ser Ser
 130 135 140

(2) INFORMATION FOR SEQ ID NO:13:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 393 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 1..393

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

ATG GGA TGG AGC TGT ATC ATC CTC TTC TTG GTA GCA ACA GCT ACA GGT	48
Met Gly Trp Ser Cys Ile Ile Leu Phe Leu Val Ala Thr Ala Thr Gly	
1 5 10 15	
GTC CAC TCC GAT ATC GTG ATG ACC CAG TCT CCA GAC TCG CTA GCT GTG	96
Val His Ser Asp Ile Val Met Thr Gln Ser Pro Asp Ser Leu Ala Val	
20 25 30	
TCT CTG GGC GAG AGG GCC ACC ATC AAC TGC AAG GCC TCC CAA AGT GTT	144
Ser Leu Gly Glu Arg Ala Thr Ile Asn Cys Lys Ala Ser Gln Ser Val	
35 40 45	
GAT TAT GAT GGT GAT AGT TAT ATG AAC TGG TAT CAG CAG AAA CCC GGG	192
Asp Tyr Asp Gly Asp Ser Tyr Met Asn Trp Tyr Gln Gln Lys Pro Gly	
50 55 60	
CAG CCT CCT AAG TTG CTC ATT TAC GCT GCA TCC AAT CTA GAA TCT GGG	240
Gln Pro Pro Lys Leu Leu Ile Tyr Ala Ala Ser Asn Leu Glu Ser Gly	
65 70 75 80	
GTA CCT GAC CGA TTC AGT GGC AGC GGG TCT GGG ACA GAT TTC ACT CTC	288
Val Pro Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu	
85 90 95	
ACC ATC AGC AGC CTG CAG GCT GAA GAT GTG GCA GTA TAC TAC TGT CAG	336
Thr Ile Ser Ser Leu Gln Ala Glu Asp Val Ala Val Tyr Tyr Cys Gln	

100	105	110	
CAA AGT AAT GAG GAT CCT CCG AGG TTC GGC GGA GGG ACC AAG GTG GAG			384
Gln Ser Asn Glu Asp Pro Pro Arg Phe Gly Gly Gly Thr Lys Val Glu			
115	120	125	
ATC AAA CGT			393
Ile Lys Arg			
130			

(2) INFORMATION FOR SEQ ID NO:14:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 131 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

Met Gly Trp Ser Cys Ile Ile Leu Phe Leu Val Ala Thr Ala Thr Gly			
1	5	10	15
Val His Ser Asp Ile Val Met Thr Gln Ser Pro Asp Ser Leu Ala Val			
20	25	30	
Ser Leu Gly Glu Arg Ala Thr Ile Asn Cys Lys Ala Ser Gln Ser Val			
35	40	45	
Asp Tyr Asp Gly Asp Ser Tyr Met Asn Trp Tyr Gln Gln Lys Pro Gly			
50	55	60	
Gln Pro Pro Lys Leu Leu Ile Tyr Ala Ala Ser Asn Leu Glu Ser Gly			
65	70	75	80

Val Pro Asp Arg Phe Ser Gly Ser Gly Thr Asp Phe Thr Leu
 85 90 95

Thr Ile Ser Ser Leu Gln Ala Glu Asp Val Ala Val Tyr Tyr Cys Gln
 100 105 110

Gln Ser Asn Glu Asp Pro Pro Arg Phe Gly Gly Gly Thr Lys Val Glu
 115 120 125

Ile Lys Arg
 130

(2) INFORMATION FOR SEQ ID NO:15:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 45 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 1..45

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

AAG GCC AGC CAA AGT GTT GAT TAT GAT GGT GAT AGT TAT ATG AAC 45
 Lys Ala Ser Gln Ser Val Asp Tyr Asp Gly Asp Ser Tyr Met Asn
 1 5 10 15

(2) INFORMATION FOR SEQ ID NO:16:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 15 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

Lys Ala Ser Gln Ser Val Asp Tyr Asp Gly Asp Ser Tyr Met Asn
1 5 10 15

(2) INFORMATION FOR SEQ ID NO:17:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 21 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 1..21

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

GCT GCA TCC AAT CTA GAA TCT
Ala Ala Ser Asn Leu Glu Ser
1 5

21

(2) INFORMATION FOR SEQ ID NO:18:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 7 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

Ala Ala Ser Asn Leu Glu Ser

1 5

(2) INFORMATION FOR SEQ ID NO:19:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 27 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: double

(D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 1..27

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

CAG CAA AGT AAT GAG GAT CCT CCG ACG

27

Gln Gln Ser Asn Glu Asp Pro Pro Thr

1 5

(2) INFORMATION FOR SEQ ID NO:20:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 9 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

Gln Gln Ser Asn Glu Asp Pro Pro Thr
1 5

(2) INFORMATION FOR SEQ ID NO:21:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 21 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 1..21

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

ACT TCT GGT ATG GGT GTG AGC
Thr Ser Gly Met Gly Val Ser
1 5

21

(2) INFORMATION FOR SEQ ID NO:22:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 7 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

Thr Ser Gly Met Gly Val Ser
1 5

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 33 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: cDNA

- (ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION: 1..33

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

AGA GAG ACT GTG TTC TAC TGG TAC TTC GAT GTC
Arg Glu Thr Val Phe Tyr Trp Tyr Phe Asp Val
1 5 10

33

(2) INFORMATION FOR SEQ ID NO:26:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 11 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

Arg Glu Thr Val Phe Tyr Trp Tyr Phe Asp Val
1 5 10

(2) INFORMATION FOR SEQ ID NO:27:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 27 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 1..27

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:

CAG CAA AGT AAT GAG GAT CCT CCG AGG
Gln Gln Ser Asn Glu Asp Pro Pro Arg
1 5

27

(2) INFORMATION FOR SEQ ID NO:28:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 9 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:

Gln Gln Ser Asn Glu Asp Pro Pro Arg
1 5

(2) INFORMATION FOR SEQ ID NO:29:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 36 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:29:

CTAACTCA TTCCTGTTGA AGCTCTTGAC AATGGG

36

(2) INFORMATION FOR SEQ ID NO:30:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 29 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:30:

GTACATATGC AAGGCTTACA ACCACAATC

29

(2) INFORMATION FOR SEQ ID NO:31:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 117 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:31:

GGTTACCCTG CGTGAATCCG GTCCGGCACT AGTTAAACCG ACCCAGACCC TGACGTTAAC 60
CTGCACCTTC TCCGGTTTCT CCCTGTCGAC CTCCGGTATG GGTGTTTCCT GGATCCG 117

(2) INFORMATION FOR SEQ ID NO:32:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 120 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:32:

TCAGCCGCCG GGTAAGGTC TAGAATGGCT GGCTCACATC TACTGGGACG ACGACAAACG 60
TTACAACCCG AGCCTGAAAT CCCGTCTGAC GATATCCAAA GACACCTCCC GTAACCAGGT 120

(2) INFORMATION FOR SEQ ID NO:33:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 120 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:33:

TGTTCTGACC ATGGACCCGG TTGACACCGC TACCTACTAC TGCGCTCGTC GCGAAACCGT 60

TTTCTACTGG TACTTCGACG TTTGGGGTCG TGGTACCCCA GTTACCGTGA GCTCCCAACC 120

(2) INFORMATION FOR SEQ ID NO:34:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 25 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:34:

ACCCGGCGGC TGACGGATCC AGGAA 25

(2) INFORMATION FOR SEQ ID NO:35:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:35:

ATGGTCAGAA CAACCTGGTT ACGG 24

(2) INFORMATION FOR SEQ ID NO:36:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 25 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:36:

TTCGGGTTAC CCTGCGTGAA TCCGG

25

(2) INFORMATION FOR SEQ ID NO:37:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 21 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:37:

CCAACCCTCG AGTGCCATTG A

21

(2) INFORMATION FOR SEQ ID NO:38:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 43 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:38:

CTAGCTGTGT CTCTGGGCGA GAGGGCCACC ATCAACTGCA AGG

43

(2) INFORMATION FOR SEQ ID NO:39:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 39 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:39:

CCTTGCAGTT GATGGTGGCC CTCTCGCCCA GAGACACAG

39

(2) INFORMATION FOR SEQ ID NO:40:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 67 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:40:

TCGAGAGGCC TCCCAAAGTG TTGATTATGA TGGTGATAGT TATATGAACT GGTATCAGCA

60

GAAACCC

67

(2) INFORMATION FOR SEQ ID NO:41:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 63 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:41:

GGGTTTCTGC TGATACCAGT TCATATAACT ATCACCATCA TAATCAACAC TTTGGGAGGC 60
CTC 63

(2) INFORMATION FOR SEQ ID NO:42:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 51 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:42:

ATACTACTGT CAGCAAAGTA ATGAGGATCC TCCGAGGTTT GCGGAGGGA C 51

(2) INFORMATION FOR SEQ ID NO:43:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 53 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:43:

CTTGGTCCCT CCGCCGAACC TCGGAGGATC CTCATTACTT TGCTGACAGT AGT

53

(2) INFORMATION FOR SEQ ID NO:44:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 55 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:44:

GGGCAGCCTC CTAAGTTGCT CATTACGCT GCATCCAATC TAGAATCTGG GGTAC

55

(2) INFORMATION FOR SEQ ID NO:45:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 51 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:45:

CCCAGATTCT AGATTGGATG CAGCGTAAAT GAGCAACTTA GGAGGCTGCC C

51

(2) INFORMATION FOR SEQ ID NO:46:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 83 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:46:

AATTCGAGGA CGCCAGCAAC ATGGTGTTGC AGACCCAGGT CTCATTCTCT CTGTTGCTCT 60
GGATCTCTGG TGCCTACGGG CAG 83

(2) INFORMATION FOR SEQ ID NO:47:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 84 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:47:

GTAACCTGCC CGTAGGCACC AGAGATCCAG AGCAACAGAG AAATGAAGAC CTGGGTCTGC 60
AACACCATGT TGCTGGCGTC CTCG 84

(2) INFORMATION FOR SEQ ID NO:48:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:48:

CAGGTTACCC TGAAAGAGTC

20

(2) INFORMATION FOR SEQ ID NO:49:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 18 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:49:

GAAGTAGTCC TTGACCAG

18

(2) INFORMATION FOR SEQ ID NO:50:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 31 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:50:

GTCACCGTCT CCTCAGCTAG CACCAAGGGG C

31

(2) INFORMATION FOR SEQ ID NO:51:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 22 base pairs

- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:51:

CTTGGTGCTA GCTGAGGAGA CG

22

(2) INFORMATION FOR SEQ ID NO:52:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 47 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:52:

CATCTAGATG GCGCCGCCAC AGTACGTTTG ATCTCCAGCT TGGTCCC

47

(2) INFORMATION FOR SEQ ID NO:53:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 45 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: double
 - (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:53:

AAGGCCTCCC AAAGTGTGA TTATGATGGT GATAGTTATA TGAAC

45

(2) INFORMATION FOR SEQ ID NO:54:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 21 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:54:

ACCTCCGGTA TGGGTGTTTC C

21

(2) INFORMATION FOR SEQ ID NO:55:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 48 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:55:

CACATCTACT GGGACGACGA CAAACGTTAC AACCCGAGCC TGAAATCC

48

(2) INFORMATION FOR SEQ ID NO:56:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 33 base pairs

- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:56:

CGCGAAACCG TTTCTACTG GTACTTCGAC GTT

33

(2) INFORMATION FOR SEQ ID NO:57:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 393 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 1..393

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:57:

ATG GGA TGG AGC TGT ATC ATC CTC TTC TTG GTA GCA ACA GCT ACA GGT	48
Met Gly Trp Ser Cys Ile Ile Leu Phe Leu Val Ala Thr Ala Thr Gly	
1 5 10 15	
GTC CAC TCC GAT ATC GTG ATG ACC CAG TCT CCA GAC TCG CTA GCT GTG	96
Val His Ser Asp Ile Val Met Thr Gln Ser Pro Asp Ser Leu Ala Val	
20 25 30	
TCT CTG GGC GAG AGG GCC ACC ATC AAC TGC AAG GCC TCC CAA AGT GTT	144
Ser Leu Gly Glu Arg Ala Thr Ile Asn Cys Lys Ala Ser Gln Ser Val	
35 40 45	

GAT TAT GAT GGT GAT AGT TAT ATG AAC TGG TAT CAG CAG AAA CCC GGG	192
Asp Tyr Asp Gly Asp Ser Tyr Met Asn Trp Tyr Gln Gln Lys Pro Gly	
50 55 60	
CAG CCT CCT AAG TTG CTC ATT TAC GCT GCA TCC AAT CTA GAA TCT GGG	240
Gln Pro Pro Lys Leu Leu Ile Tyr Ala Ala Ser Asn Leu Glu Ser Gly	
65 70 75 80	
GTA CCT GAC CGA TTC AGT GGC AGC GGG TCT GGG ACA GAT TTC ACT CTC	288
Val Pro Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu	
85 90 95	
ACC ATC AGC AGC CTG CAG GCT GAA GAT GTG GCA GTA TAC TAC TGT CAG	336
Thr Ile Ser Ser Leu Gln Ala Glu Asp Val Ala Val Tyr Tyr Cys Gln	
100 105 110	
CAA AGT AAT GAG GAT CCT CCG ACG TTC GGC GGA GGG ACC AAA GTG GAG	384
Gln Ser Asn Glu Asp Pro Pro Thr Phe Gly Gly Gly Thr Lys Val Glu	
115 120 125	
ATC AAA CGT	393
Ile Lys Arg	
130	

(2) INFORMATION FOR SEQ ID NO:58:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 131 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:58:

Met Gly Trp Ser Cys Ile Ile Leu Phe Leu Val Ala Thr Ala Thr Gly
1 5 10 15

Val His Ser Asp Ile Val Met Thr Gln Ser Pro Asp Ser Leu Ala Val
20 25 30

Ser Leu Gly Glu Arg Ala Thr Ile Asn Cys Lys Ala Ser Gln Ser Val
35 40 45

Asp Tyr Asp Gly Asp Ser Tyr Met Asn Trp Tyr Gln Gln Lys Pro Gly
50 55 60

Gln Pro Pro Lys Leu Leu Ile Tyr Ala Ala Ser Asn Leu Glu Ser Gly
65 70 75 80

Val Pro Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu
85 90 95

Thr Ile Ser Ser Leu Gln Ala Glu Asp Val Ala Val Tyr Tyr Cys Gln
100 105 110

Gln Ser Asn Glu Asp Pro Pro Thr Phe Gly Gly Gly Thr Lys Val Glu
115 120 125


Ile Lys Arg
130

Applicant's or agent's file reference number P50186-2	International Application No. PCT/US94/10308
---	--

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13*bis*)

A. The indications made below relate to the microorganism referred to in the description on page 32, line 14.	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution European Collection of Animal Cell Cultures (ECACC)	
Address of depositary institution (including postal code and country) Porton Down Salisbury Wiltshire, SP4 0JG United Kingdom	
Date of Deposit 06 October 1993	Accession Number 93100620
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
In respect of those designations in which a European or Australian Patent is sought or in any other states having equivalent provisions, a sample of the deposited micro-organism will be made available until the publication of the mention of the grant of the patent or until the date on which the application has been refused or withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample.	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States) All	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications, e.g., "Accession Number of Deposit")	

For receiving Office use only	For International Bureau use only
<input checked="" type="checkbox"/> This sheet was received with the international application	<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer 	Authorized officer

WHAT IS CLAIMED IS:

1. A fusion protein having binding specificity for human interleukin-4 (IL4) which comprises complementarity determining regions (CDRs) derived from a non-human neutralizing monoclonal antibody characterized by a dissociation constant equal to or less than 2×10^{-10} M for human IL4, and a first fusion partner.
2. The fusion protein according to claim 1 which is operatively linked to a second fusion partner.
3. The fusion protein according to claim 1 wherein said non-human neutralizing monoclonal antibody is selected from the group consisting of 3B9 and 6A1.
4. The fusion protein according to claim 2 wherein said second fusion partner comprises all or part of an immunoglobulin constant heavy chain or immunoglobulin constant light chain, or both.
5. The fusion protein according to claim 1 wherein said first fusion partner sequence is the heavy chain sequence of: amino acids 21-50, 56-71, 88-119, and 131-141 of SEQ ID NO:12.
6. The fusion protein according to claim 1 wherein said first fusion partner sequence is the light chain sequence of: amino acids 20-42, 58-72, 80-111, and 121-131 of SEQ ID NO: 14.
7. The fusion protein according to claim 1 wherein said amino acid sequences of the complementarity determining regions for the heavy chain are:
 - (a) ThrSerGlyMetGlyValSer: SEQ ID NO:22,
 - (b) HisIleTyrTrpAspAspAspLysArgTyrAsnPro-SerLeuLysSer: SEQ ID NO:24, or
 - (c) ArgGluThrValPheTyrTrpPheAspVal: SEQ ID NO:26.
8. The fusion protein according to claim 1 wherein said amino acid sequences of the complementarity determining regions for the light chain are:
 - (a) LeuAlaSerGlnSerValAspTyrAspGlyAspSerTyrMetAsn: SEQ ID NO:16,

- (b) AlaAlaSerAsnLeuGluSer: SEQ ID NO:18, or
- (c) GlnGlnSerAsnGluAspProProArg: SEQ ID NO:28.

9. The fusion protein according to claim 1 wherein said amino acid
 5 sequences of the complementarity determining regions for the light chain are:

- (a) LysAlaSerGlnSerValAspTyrAspGlyAspSerTyrMetAsn: SEQ ID
 NO:16,
- (b) AlaAlaSerAsnLeuGluSer: SEQ ID NO:18, or
- (c) GlnGlnSerAsnGluAspProProThr: SEQ ID NO:20.

10

10. An immunoglobulin heavy chain complementarity determining
 region (CDR), the amino acid sequence of which is selected from the group
 consisting of:

- (a) ThrSerGlyMetGlyValSer: SEQ ID NO:22,
- 15 (b) HisIleTyrTrpAspAspAspLysArgTyrAsnPro-
 SerLeuLysSer: SEQ ID NO:24, and
- (c) ArgGluThrValPheTyrTrpPheAspVal: SEQ ID NO:26.

11. An immunoglobulin light chain complementarity determining region
 20 (CDR), the amino acid sequence of which is selected from the group consisting of:

- (a) LeuAlaSerGlnSerValAspTyrAspGlyAspSerTyrMetAsn: SEQ ID
 NO:16,
- (b) AlaAlaSerAsnLeuGluSer: SEQ ID NO:18,
- (c) GlnGlnSerAsnGluAspProProArg: SEQ ID NO:28; and
- 25 (d) GlnGlnSerAsnGluAspProProThr: SEQ ID NO:20.

12. A nucleic acid molecule encoding an immunoglobulin heavy chain
 complementarity determining region (CDR), the sequence of which is selected from
 the group consisting of:

- 30 (a) ACT TCT GGT ATG GGT GTG AGC: SEQ ID NO:21,
- (b) CAC ATT TAC TGG GAT GAT GAC AAG CGC TAT
 AACCCATCCCTGAAGAGC: SEQ ID NO:23,
- (c) AGA GAG ACT GTG TTC TAC TGG TAC TTC GAT
 GTC: SEQ ID NO:25,
- 35 (d) ACC TCC GGT ATG GGT GTT TCC: SEQ ID NO: 54,
- (e) CAC ATC TAC TGG GAC GAC GAC AAA CGT TAC AAC CCG
 AGC CTG AAA TCC: SEQ ID NO: 55, and

(f) CGC GAA ACC GTT TTC TAC TGG TAC TTC GAC GTT: SEQ
ID NO: 56.

13. A nucleic acid molecule encoding an immunoglobulin light chain
5 complementarity determining region (CDR), the sequence of which is selected from
the group consisting of:

(a) AAG GCC AGC CAA AGT GTT GAT TAT GAT GGT
GAT AGT TAT ATG AAC: SEQ ID NO:15,

(b) AAG GCC TCC CAA AGT GTT GAT TAT GAT GGT GAT AGT
10 TAT ATG AAC: SEQ ID NO: 53,

(c) GCT GCA TCC AAT CTA GAA TCT: SEQ ID NO:17, (d)
CAG CAA AGT AAT GAG GAT CCT CCG ACG: SEQ ID NO:19, and

(e) CAG CAA AGT AAT GAG GAT CCT CCG AGG: SEQ ID NO:27.

14. A humanized antibody comprising a heavy chain and a light chain,
15 said antibody characterized by a dissociation constant equal to or less than about 2×10^{-10} M for human IL4, wherein the framework regions of said heavy and light
chains are derived from at least one selected human antibody and the amino acid
sequences of the complementarity determining regions of each said chain are
20 derived from a non-human neutralizing monoclonal antibody specific for human IL4
characterized by a dissociation constant equal to or less than about 2×10^{-10} M for
human IL4.

15. The antibody according to claim 14 wherein said antibody is
25 optionally linked to an effector agent selected from the group consisting of a non-
protein carrier molecule, polystyrene, and plastic beads.

16. A chimeric antibody comprising a heavy chain and a light chain, said
antibody characterized by a dissociation constant equal to or less than about 2×10^{-10}
30 M for human IL4, wherein the amino acid sequences of the complementarity
determining regions of said heavy chain and said light chain are derived from a non-
human neutralizing monoclonal antibody specific for human IL4 characterized by a
dissociation constant equal to or less than about 2×10^{-10} M for human IL4.

17. A pharmaceutical composition comprising the fusion protein of claim
35 1 and a pharmaceutically acceptable carrier.

18. A method of treating allergies and other conditions associated with excess IgE production in a human comprising the step of administering to said human in need thereof an effective amount of the fusion protein of claim 1.

5 19. An isolated nucleic acid sequence which is selected from the group consisting of:

- (a) a nucleic acid sequence encoding the fusion protein of claim 1;
 - (b) a nucleic acid sequence complementary to (a);
 - 10 (c) a nucleic acid sequence of 18 or more nucleotides capable of hybridizing to (a) or (b) under stringent conditions; and
 - (d) a fragment or analog of (a), (b), or (c) which encodes a protein characterized by having specificity for human interleukin-4;
- wherein said sequence optionally contains a restriction site.

15

20. The isolated nucleic acid sequence according to claim 19, wherein the sequence encoding the fusion protein comprises the nucleic acid sequence of Fig. 5, SEQ ID NO:13.

20 21. The isolated nucleic acid sequence according to claim 19, wherein the sequence encoding the fusion protein comprises the nucleic acid sequence of Fig. 4, SEQ ID NO:11.

25 22. An isolated nucleic acid sequence which is selected from the group consisting of:

- (a) a nucleic acid sequence encoding a complementarity determining region (CDR) wherein said CDR is obtained from a neutralizing murine monoclonal antibody specific for human interleukin-4 and having a dissociation constant equal to or less than about 2×10^{-10} M;
- 30 (b) a nucleic acid sequence complementary to (a);
- (c) a nucleic acid sequence of 18 or more nucleotides capable of hybridizing under stringent conditions to (a) or (b); and
- (d) a fragment or analog of (a), (b) or (c) which encodes a protein characterized by having specificity for human interleukin-4.

35

23. The isolated nucleic acid sequence according to claim 22, wherein said sequence is selected from the group of heavy chain complementarity determining region-encoding sequences consisting of:

- (a) ACT TCT GGT ATG GGT GTG AGC: SEQ ID NO:21,
- 5 (b) CAC ATT TAC TGG GAT GAT GAC AAG CGC TAT
AAC CCA TCC CTG AAG AGC: SEQ ID NO:23,
- (c) AGA GAG ACT GTG TTC TAC TGG TAC TTC GAT
GTC: SEQ ID NO:25,
- (d) ACC TCC GGT ATG GGT GTT TCC: SEQ ID NO: 54,
- 10 (e) CAC ATC TAC TGG GAC GAC GAC AAA CGT TAC AAC CCG
AGC CTG AAA TCC: SEQ ID NO: 55, and
- (f) CGC GAA ACC GTT TTC TAC TGG TAC TTC GAC GTT:
SEQ ID NO: 56.

24. The isolated nucleic acid sequence according to claim 22, wherein said sequence is selected from the group of light chain complementarity determining region-encoding sequences consisting of:

- (a) AAG GCC AGC CAA AGT GTT GAT TAT GAT GGT
GAT AGT TAT ATG AAC: SEQ ID NO:15,
- 20 (b) AAG GCC TCC CAA AGT GTT GAT TAT GAT GGT GAT AGT
TAT ATG AAC: SEQ ID NO: 53,
- (c) GCT GCA TCC AAT CTA GAA TCT: SEQ ID NO:17, (d)
CAG CAA AGT AAT GAG GAT CCT CCG ACG: SEQ ID NO:19, and
- (e) CAG CAA AGT AAT GAG GAT CCT CCG AGG: SEQ ID NO:27.

25. A recombinant plasmid comprising the nucleic acid sequence of claim 19.

26. A recombinant plasmid comprising the nucleic acid sequence of claim 22.

27. A host cell transfected with the recombinant plasmid of claim 25.

28. A host cell transfected with the recombinant plasmid of claim 26.

29. A process for producing a humanized antibody specific for human interleukin-4 comprising culturing a cell line transfected with the recombinant

plasmid of claim 25 under the control of selected regulatory sequences capable of directing the expression thereof in said cells.

30. A method for diagnosing allergies and other conditions associated with excess immunoglobulin E production in a human which comprises contacting a sample of biological fluid with a high titer monoclonal antibody for human IL4 and assaying for the occurrence of binding between said monoclonal antibody and human interleukin 4.

31. A method for screening monoclonal antibodies which have a high titer for human interleukin 4 which comprises:

- a) preparing a hybridoma cell line characterized by secretion of a monoclonal antibody to human interleukin 4; and
- b) screening said hybridoma cell line with aldehyde-coupled human interleukin-4 or biotinylated human interleukin-4.

32. A neutralizing monoclonal antibody having a high titer for human interleukin-4, a Fab fragment or a F(ab')₂ fragment thereof, produced by screening a library of hybridoma products with aldehyde-coupled human interleukin-4 or biotinylated human interleukin-4.

33. A rodent neutralizing monoclonal antibody specific for human interleukin-4 and having a binding affinity characterized by a dissociation constant equal to or less than about 2×10^{-10} M.

34. The monoclonal antibody according to claim 33 wherein said rodent is a mouse.

35. The monoclonal antibody according to claim 34, which comprises the light chain amino acid sequence of SEQ ID NO: 2, and the heavy chain amino acid sequence of SEQ ID NO: 4.

36. The monoclonal antibody according to claim 33, wherein said rodent is a rat.

37. The monoclonal antibody according to claim 36 having the identifying characteristics of 6A1.

38. A hybridoma having the identifying characteristics of cell line 3426A11C1B9.

FIGURE 1

Murine Antibody 3B9 Light Chain
Native Signal Sequence and Variable Region

Nucleotide Sequence SEQ ID NO:1

Amino Acid Sequence SEQ ID NO:2

ATG GAG ACA GAC ACA ATC CTG CTA TGG GTG CTG CTG CTC	39
Met Glu Thr Asp Thr Ile Leu Leu Trp Val Leu Leu Leu	
1 5 10	
TGG GTT CCA GGC TCC ACT GGT GAC ATT GTG CTG ACC CAA	78
Trp Val Pro Gly Ser Thr Gly Asp Ile Val Leu Thr Gln	
15 20 25	
TCT CCA GCT TCT TTG GCT GTG TCT CTA GGG CAG AGG GCC	117
Ser Pro Ala Ser Leu Ala Val Ser Leu Gly Gln Arg Ala	
30 35	
ACC ATC TCC TGC AAG GCC AGC CAA AGT GTT GAT TAT GAT	156
Thr Ile Ser Cys Lys Ala Ser Gln Ser Val Asp Tyr Asp	
40 45 50	
GGT GAT AGT TAT ATG AAC TGG TAC CAA CAG AAA CCA GGA	195
Gly Asp Ser Tyr Met Asn Trp Tyr Gln Gln Lys Pro Gly	
55 60 65	
CAG CCA CCC AAA CTC CTC ATC TAT GCT GCA TCC AAT CTA	234
Gln Pro Pro Lys Leu Leu Ile Tyr Ala Ala Ser Asn Leu	
70 75	
GAA TCT GGG ATC CCA GCC AGG TTT AGT GGC AGT GGG TCT	273
Glu Ser Gly Ile Pro Ala Arg Phe Ser Gly Ser Gly Ser	
80 85 90	
GGG ACA GAC TTC ACC CTC AAC ATC CAT CCT GTG GAG GAG	312
Gly Thr Asp Phe Thr Leu Asn Ile His Pro Val Glu Glu	
95 100	
GAG GAT GCT GCA ACC TAT TAC TGT CAG CAA AGT AAT GAG	351
Glu Asp Ala Ala Thr Tyr Tyr Cys Gln Gln Ser Asn Glu	
105 110 115	
GAT CCT CCG ACG TTC GGT GGA GGC ACC AAG CTG GAA ATC	390
Asp Pro Pro Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile	
120 125 130	
AAA CGG	396
Lys Arg	

Figure 2

Murine Antibody 3B9 Heavy Chain
Native Signal Sequence and Variable Region

Nucleotide Sequence SEQ ID NO: 3

Amino Acid Sequence SEQ ID NO: 4

GAATTCGCGG CCGCTATGCA GGGACAATCA GCAGCAGCAA	40
TGAGGAAGTA AGCCTGTGCA GAT ATG AAC AGG CTT ACT TCC	81
Met Asn Arg Leu Thr Ser	
1 5	
TCA TTG CTG CTG CTG ATT GTC CCT GCA TAT GTC CTG TCC	120
Ser Leu Leu Leu Leu Ile Val Pro Ala Tyr Val Leu Ser	
10 15	
CAG GTT ACT CTG AAA GAG TCT GGC CCT GGG ATA TTG CAG	159
Gln Val Thr Leu Lys Glu Ser Gly Pro Gly Ile Leu Gln	
20 25 30	
CCC TCC CAG ACC CTC AGT CTG ACT TGT TCT TTC TCT GGG	198
Pro Ser Gln Thr Leu Ser Leu Thr Cys Ser Phe Ser Gly	
35 40 45	
TTT TCA CTG AGC ACT TCT GGT ATG GGT GTG AGC TGG ATT	237
Phe Ser Leu Ser Thr Ser Gly Met Gly Val Ser Trp Ile	
50 55	
CGT CAG CCT TCA GGA AAG GGT CTG GAG TGG CTG GCA CAC	276
Arg Gln Pro Ser Gly Lys Gly Leu Glu Trp Leu Ala His	
60 65 70	
ATT TAC TGG GAT GAT GAC AAG CGC TAT AAC CCA TCC CTG	315
Ile Tyr Trp Asp Asp Asp Lys Arg Tyr Asn Pro Ser Leu	
75 80	
AAG AGC CGG CTC ACA ATC TCC AAG GAT ACC TCC AGC AAC	354
Lys Ser Arg Leu Thr Ile Ser Lys Asp Thr Ser Ser Asn	
85 90 95	
CAG GTA TTC CTC AAG ATC ACC AGT GTG GAC ACT GCA GAT	393
Gln Val Phe Leu Lys Ile Thr Ser Val Asp Thr Ala Asp	
100 105 110	
ACT GCC ACA TAC TAC TGT GCT CGA AGA GAG ACT GTG TTC	432
Thr Ala Thr Tyr Tyr Cys Ala Arg Arg Glu Thr Val Phe	
115 120	

Figure 2 (con't)

TAC	TGG	TAC	TTC	GAT	GTC	TGG	GGC	GCA	GGG	ACC	ACG	GTC	471
<u>Tyr</u>	<u>Trp</u>	<u>Tyr</u>	<u>Phe</u>	<u>Asp</u>	<u>Val</u>	Trp	Gly	Ala	Gly	Thr	Thr	Val	
125						130					135		
ACC	GTC	TCC	TCA										483
Thr	Val	Ser	Ser										
			140										

FIGURE 3

Human/Murine 3B9 Chimeric Antibody Heavy Chain
Signal Sequence and Variable Region

Nucleotide Sequence SEQ ID NO: 9

Amino Acid Sequence SEQ ID NO:10

ATG GTG TTG CAG ACC CAG GTC TTC ATT TCT CTG TTG CTC	39
Met Val Leu Gln Thr Gln Val Phe Ile Ser Leu Leu Leu	
1 5 10	
TGG ATC TCT GGT GCC TAC GGG CAG GTT ACC CTG AAA GAG	78
Trp Ile Ser Gly Ala Tyr Gly Gln Val Thr Leu Lys Glu	
15 20 25	
TCT GGC CCT GGG ATA TTG CAG CCC TCC CAG ACC CTC AGT	117
Ser Gly Pro Gly Ile Leu Gln Pro Ser Gln Thr Leu Ser	
30 35	
CTG ACT TGT TCT TTC TCT GGG TTT TCA CTG AGC ACT TCT	156
Leu Thr Cys Ser Phe Ser Gly Phe Ser Leu Ser <u>Thr Ser</u>	
40 45 50	
GGT ATG GGT GTG AGC TGG ATT CGT CAG CCT TCA GGA AAG	195
<u>Gly Met Gly Val Ser</u> Trp Ile Arg Gln Pro Ser Gly Lys	
55 60 65	
GGT CTG GAG TGG CTG GCA CAC ATT TAC TGG GAT GAT GAC	234
Gly Leu Glu Trp Leu Ala <u>His Ile Tyr Trp Asp Asp Asp</u>	
70 75	
AAG CGC TAT AAC CCA TCC CTG AAG AGC CGG CTC ACA ATC	273
<u>Lys Arg Tyr Asn Pro Ser Leu Lys Ser</u> Arg Leu Thr Ile	
80 85 90	
TCC AAG GAT ACC TCC AGC AAC CAG GTA TTC CTC AAG ATC	312
Ser Lys Asp Thr Ser Ser Asn Gln Val Phe Leu Lys Ile	
95 100	
ACC AGT GTG GAC ACT GCA GAT ACT GCC ACA TAC TAC TGT	351
Thr Ser Val Asp Thr Ala Asp Thr Ala Thr Tyr Tyr Cys	
105 110 115	
GCT CGA AGA GAG ACT GTG TTC TAC TGG TAC TTC GAT GTC	390
Ala Arg <u>Arg Glu Thr Val Phe Tyr Trp Tyr Phe Asp Val</u>	
120 125 130	
TGG GGC GCA GGG ACC ACG GTC ACC GTC TCC TCA	423
Trp Gly Ala Gly Thr Thr Val Thr Val Ser Ser	
135 140	

FIGURE 4

Humanized 3B9 Antibody Heavy Chain
Signal Sequence and Variable Region

Nucleotide Sequence SEQ ID NO: 11

Amino Acid Sequence SEQ ID NO: 12

ATG GTG TTG CAG ACC CAG GTC TTC ATT TCT CTG TTG CTC	39
Met Val Leu Gln Thr Gln Val Phe Ile Ser Leu Leu Leu	
1 5 10	
TGG ATC TCT GGT GCC TAC GGG CAG GTT ACC CTG CGT GAA	78
Trp Ile Ser Gly Ala Tyr Gly Gln Val Thr Leu Arg Glu	
15 20 25	
TCC GGT CCG GCA CTA GTT AAA CCG ACC CAG ACC CTG ACG	117
Ser Gly Pro Ala Leu Val Lys Pro Thr Gln Thr Leu Thr	
30 35	
TTA ACC TGC ACC TTC TCC GGT TTC TCC CTG TCG ACC TCC	156
Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Ser	
40 45 50	
GGT ATG GGT GTT TCC TGG ATC CGT CAG CCG CCG GGT AAA	195
Gly Met Gly Val Ser Trp Ile Arg Gln Pro Pro Gly Lys	
55 60 65	
GGT CTA GAA TGG CTG GCT CAC ATC TAC TGG GAC GAC GAC	234
Gly Leu Glu Trp Leu Ala His Ile Tyr Trp Asp Asp Asp	
70 75	
AAA CGT TAC AAC CCG AGC CTG AAA TCC CGT CTG ACG ATA	273
Lys Arg Tyr Asn Pro Ser Leu Lys Ser Arg Leu Thr Ile	
80 85 90	
TCC AAA GAC ACC TCC CGT AAC CAG GTT GTT CTG ACC ATG	312
Ser Lys Asp Thr Ser Arg Asn Gln Val Val Leu Thr Met	
95 100	
ACT AAC ATG GAC CCG GTT GAC ACC GCT ACC TAC TAC TGC	351
Thr Asn Met Asp Pro Val Asp Thr Ala Thr Tyr Tyr Cys	
105 110 115	
GCT CGA CGC GAA ACC GTT TTC TAC TGG TAC TTC GAC GTT	390
Ala Arg Arg Glu Thr Val Phe Tyr Trp Tyr Phe Asp Val	
120 125 130	
TGG GGT CGT GGT ACC CCA GTT ACC GTG AGC TCA	423
Trp Gly Arg Gly Thr Pro Val Thr Val Ser Ser	
135 140	

FIGURE 5

Humanized 3B9 Antibody Light Chain
Signal Sequence and Variable Region

Nucleotide Sequence SEQ ID NO: 13

Amino Acid Sequence SEQ ID NO: 14

ATG GGA TGG AGC TGT ATC ATC CTC TTC TTG GTA GCA ACA	39
Met Gly Trp Ser Cys Ile Ile Leu Phe Leu Val Ala Thr	
1 5 10	
GCT ACA GGT GTC CAC TCC GAT ATC GTG ATG ACC CAG TCT	78
Ala Thr Gly Val His Ser Asp Ile Val Met Thr Gln Ser	
15 20 25	
CCA GAC TCG CTA GCT GTG TCT CTG GGC GAG AGG GCC ACC	117
Pro Asp Ser Leu Ala Val Ser Leu Gly Glu Arg Ala Thr	
30 35	
ATC AAC TGC AAG GCC TCC CAA AGT GTT GAT TAT GAT GGT	156
Ile Asn Cys Lys Ala Ser Gln Ser Val Asp Tyr Asp Gly	
40 45 50	
GAT AGT TAT ATG AAC TGG TAT CAG CAG AAA CCC GGG CAG	195
Asp Ser Tyr Met Asn Trp Tyr Gln Gln Lys Pro Gly Gln	
55 60 65	
CCT CCT AAG TTG CTC ATT TAC GCT GCA TCC AAT CTA GAA	234
Pro Pro Lys Leu Leu Ile Tyr Ala Ala Ser Asn Leu Glu	
70 75	
TCT GGG GTA CCT GAC CGA TTC AGT GGC AGC GGG TCT GGG	273
Ser Gly Val Pro Asp Arg Phe Ser Gly Ser Gly Ser Gly	
80 85 90	
ACA GAT TTC ACT CTC ACC ATC AGC AGC CTG CAG GCT GAA	312
Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Ala Glu	
95 100	
GAT GTG GCA GTA TAC TAC TGT CAG CAA AGT AAT GAG GAT	351
Asp Val Ala Val Tyr Tyr Cys Gln Gln Ser Asn Glu Asp	
105 110 115	
CCT CCG AGG TTC GGC GGA GGG ACC AAG GTG GAG ATC AAA	390
Pro Pro Arg Phe Gly Gly Gly Thr Lys Val Glu Ile Lys	
120 125 130	
CGT	393
Arg	

FIGURE 6A

Signal Sequence

Nucleotide SEQ ID NO:5

Amino Acid SEQ ID NO:6

ATG	GTG	TTG	CAG	ACC	CAG	GTC	TTC	ATT	TCT	CTG	TTG	CTC	39
Met	Val	Leu	Gln	Thr	Gln	Val	Phe	Ile	Ser	Leu	Leu	Leu	
1				5					10				

TGG	ATC	TCT	GGT	GCC	TAC								
Trp	Ile	Ser	Gly	Ala	Tyr								
	15												

FIGURE 6B

Signal Sequence

Nucleotide SEQ ID NO:7

Amino Acid SEQ ID NO:8

ATG	GGA	TGG	AGC	TGT	ATC	ATC	CTC	TTC	TTG	GTA	GCA	ACA	39
Met	Gly	Trp	Ser	Cys	Ile	Ile	Leu	Phe	Leu	Val	Ala	Thr	
1				5					10				

GCT	ACA	GGT	GTC	CAC	TCC	GAT	ATC	GTG	ATG	ACC	CAG	TCT	78
Ala	Thr	Gly	Val	His	Ser	Asp							
	15					20							

I14chlC-cdn




FIGURE 9





FIGURE 10



INTERNATIONAL SEARCH REPORT

International application No.

PCT/US94/10308

A. CLASSIFICATION OF SUBJECT MATTER IPC(6) : Please See Extra Sheet. US CL : Please See Extra Sheet. According to International Patent Classification (IPC) or to both national classification and IPC				
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) U.S. : 530/387.3, 388.25; 536/23.53; 435/7.1, 70.21, 172.2, 240.2, 240.27, 320.1; 424/133.1, 134.1, 145.1 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) APS, CAS ONLINE, BIOSIS, MEDLINE, SWISSPROT, GENESEQ 3				
C. DOCUMENTS CONSIDERED TO BE RELEVANT				
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.		
Y	WO, A, 90/07861 (QUEEN ET AL.) 26 JULY 1990, SEE ENTIRE DOCUMENT.	1-29		
Y	US, A, 5,041,381 (ABRAMS ET AL.) 20 AUGUST 1991, SEE ENTIRE DOCUMENT.	1-38		
X --- Y	JP, A, 91-187,395 (KANEKO ET AL.) 15 AUGUST 1991, SEE ENTIRE DOCUMENT.	32, 33, 36-38 ----- 1-31, 34, 35		
X --- Y	WO, A, 91/09059 (RAMANATHAN ET AL.) 27 JUNE 1991, SEE ENTIRE DOCUMENT.	32 ----- 1-31, 33-38		
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.				
<table border="0"> <tr> <td> * Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed </td> <td> "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family </td> </tr> </table>			* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family			
Date of the actual completion of the international search 09 DECEMBER 1994		Date of mailing of the international search report 09 JAN 1995		
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230		Authorized officer PAULA HUTZELL <i>P. Hutzell for</i> Telephone No. (703) 308-0196		

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US94/10308

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	JOURNAL OF IMMUNOLOGICAL METHODS, VOLUME 117, ISSUED 1989, CHRETIEN ET AL., "DEVELOPMENT OF POLYCLONAL AND MONOCLONAL ANTIBODIES FOR IMMUNOASSAY AND NEUTRALIZATION OF HUMAN INTERLEUKIN-4", PAGES 67-81, SEE ENTIRE DOCUMENT.	1-38
Y	HYBRIDOMA, VOLUME 12, NUMBER 4, ISSUED AUGUST 1993, ABE ET AL., "NOVEL IMMUNIZATION PROTOCOL AND ELISA SCREENING METHODS USED TO OBTAIN AND CHARACTERIZE MONOCLONAL ANTIBODIES SPECIFIC FOR HUMAN LIGHT CHAIN VARIABLE-REGION SUBGROUPS", PAGES 475-483, SEE ENTIRE DOCUMENT.	31, 32

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US94/10308

A. CLASSIFICATION OF SUBJECT MATTER:

IPC (6):

C07K 16/24, 16/46, 17/02; C07H 15/12; G01N 33/53; C12P 21/08, C12N 5/10, 5/20, 15/13; A61K 39/395

A. CLASSIFICATION OF SUBJECT MATTER:

US CL :

530/387.3, 388.25, 391.1; 536/23.53; 435/7.1, 70.21; 172.2, 240.2, 240.27, 320.1; 424/133.1, 134.1, 145.1